




Ministry of Health

National Guideline for the Management of Acute Malnutrition in Ethiopia



May 2019

Recommended Citation

Government of Ethiopia, Federal Ministry of Health. 2019. National Guideline for the Management of Acute Malnutrition. Addis Ababa: FMOH.

Contact Information:

Federal Ministry of Health

Post Office Box 1234

Sudan Street

Addis Ababa

Ethiopia

Telephone: +251 11 5517011

Fax: +251 11 551 9366

Email: nutritionexpert12@gmail.com or moh@ethionet.et

National Guideline for the Management of Acute Malnutrition in Ethiopia

May 2019



Foreword

Ethiopia has made tremendous strides in reducing child mortality and morbidity. According to the Demographic Health Survey, the prevalence of stunting declined from 58% in 2000 to 38% in 2016. However, the prevalence of wasting has remained fairly static at 12% in 2000 and 10% in 2016. To address malnutrition in all its forms, the Government is applying two programmatic approaches. The first focuses on increasing access to and availability of food through improved economic growth, better agricultural production systems along with promotion of good nutrition practices and prevention of malnutrition. The second approach aims to strengthen early warning systems and timely emergency response, including wide-scale delivery of services for the management of acute malnutrition.

The Federal Ministry of Health (FMOH) developed the first Protocol for the Management of Severe Acute Malnutrition (SAM) in 2007, and the Guideline for the Management of Moderate Acute Malnutrition (MAM) in 2012. This National Guideline for the Management of Acute Malnutrition in Ethiopia replaces the previous two guidelines. It includes the latest World Health Organisation (WHO) guidelines and recommendations, and emerging national and international evidence. It is also aligned to the National Nutrition Programme (NNP) II 2016-2020, the National Food and Nutrition Policy and the Health Sector Transformation Plan (HSTP) 2015/16 - 2019/20.

This National Guideline for the Management of Acute Malnutrition in Ethiopia aims to improve access to quality services for the management of SAM and MAM. Health care providers, academic institutions, development partners and individuals involved in the management of acute malnutrition should use this guideline as the primary point of reference for curative and preventive service delivery, training, monitoring and supervision.

The Federal Democratic Republic of Ethiopia, Ministry of Health, takes this opportunity to acknowledge the financial and technical support of the various partners, individuals and institutions who took part in the process of revising this guideline. Together, we shall end malnutrition in Ethiopia.

Lia Tadesse, MD, MHA
State Minister of Health



Acknowledgements

The National Guideline for the Management of Acute Malnutrition in Ethiopia has been revised to include the latest WHO guidelines and recommendations, and emerging national and international evidence on acute malnutrition. The revision process started in 2016 and was completed in 2018 after conducting a study on the implications of Ethiopia’s adoption of the 2013 WHO guidelines, particularly the admission and discharge criteria for the management of acute malnutrition.

The FMOH sincerely acknowledges the financial and technical contribution of United National International Children’s Emergency Fund (UNICEF), WHO, World Food Programme (WFP), United States Agency for International Development (USAID), Concern Worldwide, GOAL, Save the Children International (SCI), USAID Transform Primary Health Care Project, and Valid International.

Revision in 2016

The FMOH gratefully acknowledges the technical contributions of the following individuals who participated in the 2016 revision of the guideline: Israel Hailu (FMOH), Dr. Teweldebrhan Daniel (UNICEF), Kidist Negash (UNICEF), Etsegenet Asseffa (WHO), Tayech Yimer (WFP), Dr. Abdulaziz Ali (WFP), Jundi Ahmmed (WFP), Dr. Hailu Tesfaye (SCI), Molla Melese (Gonder University), Lubaba Hussein (Concern Worldwide), Mestawet Gebru (Tufts University, Africa Regional Office).

Revision in 2018

The FMOH’s Maternal and Child Health Directorate-Nutrition Case Team led the 2018 revision of the guidelines with technical and financial support of development partners.

.....

The FMOH greatly appreciates the following individuals who supported the revision and finalisation of the National Guideline for the Management of Acute Malnutrition in Ethiopia in 2018.

- ◆ Mr. Birara Melesse and Mrs. Frezer Abebe for leading and coordinating the process
- ◆ Alice Nkoroi, Acute Malnutrition Guidelines Consultant
- ◆ Members of the Community-based Management of Acute Malnutrition (CMAM) Technical Working Group (TWG): Frezer Abebe (FMOH), Dr. Yared Tadesse (FMOH), Dr. Belaynesh Yifru (FMOH), Getu Assefa (FMOH), Kebede Mengistu (FMOH), Miraf Tesfaye (FMOH), Shibru Kalbessa (FMOH), Dr. Betty Pido Lanyero (WHO), Dr. Bereket Yalew (WHO), Getahun Beyene (WHO), Halima Babyad (WHO), Jasinta Hyachits Achen (UNICEF), Nyauuma Nyasani (UNICEF), Kiyeeon Yoon (UNICEF), Banchiliyew Getahun (UNICEF), Tayech Yimer (WFP), Kemeria Barsenga (WFP), Orla Mary O’Neill [Emergency Nutrition Cluster Unit (ENCU)], Mathewos Tamiru (ENCU), Melanie Thurber (USAID), Dr. Yared Abebe (USAID Transform Primary Health Care Project), Sinksar Simeneh (SCI), Mitswat Mulaw (THDR), Nicky Dent (Concern Worldwide), Dr. Mengistu Tafesse (ABH Partners), Shiferaw Fisseha (SAKBusiness and Personal Development PLC), Yosef Beyene [Ethiopia Public Health Institute (EPHI)], Dr. Paluku Bahwere (Valid International), Helina Tufa (GOAL), Kidist Negash (HelpAge International) and Metasebia Legesse (Care International).

Finally, the FMOH would like to recognize all of the programme managers and technical staff who contributed to the finalization of the document.



Table of Contents

11	Abbreviations and Acronyms
13	Definition of Terms
17	1. Introduction
17	Purpose and Scope of the Guideline
18	Use of the Guideline
18	What is New in The Guideline
23	2. Integrating Acute Malnutrition Services into the Health System
23	2.1 Acute Malnutrition in the Context of Ethiopia
30	2.2 Roles and Responsibilities of Stakeholders
33	3. Community Outreach
33	3.1 Planning Community Outreach
34	3.2 Implementing Community Outreach
36	4. Management of Severe Acute Malnutrition (SAM)
37	4.1 Outpatient Therapeutic Programme (OTP) for the Management of SAM Without Medical Complications
47	4.2 Stabilisation Centre (SC) for the Management of SAM With Medical Complications
83	5. Management of Moderate Acute Malnutrition (MAM)
83	5.1 Principles of Supplementary Feeding Programmes (SFPs)
84	5.2 Blanket Supplementary Feeding Programme (BSFP)
85	5.3 Targeted Supplementary Feeding Programme (TSFP)
93	6. Managing Acute Malnutrition in Other Vulnerable Groups
93	6.1 Acute Malnutrition and HIV/AIDS
94	6.2 Acute Malnutrition and TB
95	6.3 Acute Malnutrition in Older People (\geq 60 Years)
96	6.4 Acute Malnutrition and Disability
98	7. Supply Management
98	7.1 Essential Supplies for Acute Malnutrition
98	7.2 Procurement
98	7.3 Requesting Supplies
100	7.4 Receiving Supplies
100	7.5 Stock Management

101	7.6 Reporting Supplies
102	8. Monitoring, Reporting and Evaluation
102	8.1 Monitoring Community Outreach
102	8.2 Monitoring Individual Patient Treatment
104	8.3 Reporting
106	8.4 Monitoring Programme Performance
109	8.5 Supportive Supervision
109	8.6 Coverage Assessment
110	References
112	Annex 1. Home Visit Checklist
113	Annex 2. Management of Acute Watery Diarrhoea (AWD) and SAM
118	Annex 3. Antibiotics for Patients with SAM
120	Annex 4. F-75 Reference Tables for Stabilisation Phase (Severe Wasting)
123	Annex 5. F-75 Reference Tables for Stabilisation Phase (Bilateral Pitting Oedema +++)
126	Annex 6. How to Insert a Nasogastric Tube (NGT)
128	Annex 7. F-100 Reference Tables
131	Annex 8. Toys for Children
132	Annex 9. F-100-Diluted Reference Table for Infant 0-6 Months Old
134	Annex 10. Decision Tree on the Management of MAM
135	Annex 11. Specialised Nutritious Foods
136	Annex 12. Nutrient Requirements for PLHIV
137	Annex 13. Essential Commodities for the Management of SAM
139	Annex 14. Essential Commodities for the Management of MAM
140	Annex 15. Bin Card
141	Annex 16. Stock Record Card
142	Annex 17. OTP Treatment and Follow-up Card
144	Annex 18. Stabilisation Centre (SC) Multi-chart
146	Annex 19. TSFP Treatment and Follow-up Card for Children 6-59 Months
147	Annex 20. TSFP Treatment and Follow-up Card for Pregnant and Lactating Women
148	Annex 21. TSFP Identification Card for Admitted Patients
149	Annex 22. Referral Slip for the Management of Acute Malnutrition
150	Annex 23. Registration Book for SAM Treatment
151	Annex 24. Registration Book for MAM Treatment - Children 6-59 Months
152	Annex 25. Registration Book for MAM Treatment – PLW
153	Annex 26. Monthly Statistics Report for SAM and MAM
154	Annex 27. Monthly Supplies Report for SAM and MAM
155	Annex 28. Acute Malnutrition Performance Monitoring Scorecard



List of Tables

26	Table 1. Assessing and Classifying Acute Malnutrition in Infants 0-6 Months
28	Table 2. Assessing and Classifying Acute Malnutrition in Children 6-59 Months
37	Table 3. OTP Admission Criteria
37	Table 4. OTP Entry Categories
38	Table 5. RUTF Appetite Test Results
39	Table 6. OTP Action Protocol
41	Table 7. Routine Medication and Treatment in OTP
42	Table 8. RUTF Reference Table (92 g Sachet Containing 500 Kcal)
45	Table 9. OTP Discharge Criteria
45	Table 10. OTP Exit Categories
47	Table 11. SC Admission Criteria
48	Table 12. SC Entry Categories
56	Table 13. Choice of Broad-Spectrum Antibiotics in SC
58	Table 14. Vitamin A Treatment Dosages
59	Table 15. Albendazole or Mebendazole
61	Table 16. Preparation of F-75 Feeds using the New Tin Formulation
63	Table 17. RUTF Reference Table (92 g Sachet Containing 500 Kcal), with the Amounts for Transition in the SC
65	Table 18. Preparation of F-100 Feeds using the New Tin Formulation
70	Table 19. Frequent Causes of Failure to Respond to Inpatient Treatment
70	Table 20. Criteria for Failure to Respond to Treatment
72	Table 21. Criteria for Discharge and Transferring Children 6-59 Months
73	Table 22. SC Exit Categories
74	Table 23. SC Admission Criteria for Infants 0-6 Months
75	Table 24. SC Entry Categories
75	Table 25. Doses of Iron Syrup if F-100 Is Used in Rehabilitation
77	Table 26. Preparation of F-100-Diluted Feeds using the New Tin Packaging
79	Table 27. Discharge Criteria for the Breastfed Infants 0-6 Months
79	Table 28. SC Exit Categories
82	Table 29. Discharge Criteria for the Non-Breastfed Infants 0-6 Months
83	Table 30. Types of SFPs

85	Table 31. TSFP Admission Criteria
85	Table 32. TSFP Entry Categories
87	Table 33. Routine Medication and Treatment for Children in TSFP
87	Table 34. Routine Medication and Treatment for PLW in TSFP
88	Table 35. FBF Ration Size per Patient
90	Table 36. RUSF Ration Size per Patient
92	Table 37. TSFP Discharge Criteria
92	Table 38. TSFP Exit Categories
95	Table 39. Admission and Discharge Criteria for SAM in Older People
96	Table 40. Admission and Discharge Criteria for MAM in Older People
99	Table 41. Example on How to Quantify Supplies Need
103	Table 42. The Filing System used in SC, OTP and TSFP
105	Table 43. Summary of Entry and Exit Categories
107	Table 44. Outcome Indicators
108	Table 45. Data Quality Assurance



List of Figures

25	Figure 1. Flow Chart on the Management of Acute Malnutrition
36	Figure 2. Decision Process for SAM Case Management
53	Figure 3. Treatment of Dehydration
55	Figure 4. Monitoring Patients on ReSoMal
77	Figure 5. Supplementary Suckling Technique
89	Figure 6. Preparation of the FBF Porridge
93	Figure 7. Cycle of Poor Nutrition and Infection
97	Figure 8. Interactions Between Malnutrition and Disability
108	Figure 9. Acute Malnutrition Data Flow



Abbreviations and Acronyms

>	Greater than	ICCM	Integrated Community Case Management
≥	Greater than or equal to	IMNCI	Integrated Management of Newborn and Childhood illness
<	Less than	IPLS	Integrated Pharmaceutical Logistics Systems
AIDS	Acquired Immune Deficiency Syndrome	IU	International Unit(s)
ART	Anti-Retroviral Therapy	IV	Intravenous
ARV	Anti-Retroviral	IYCF	Infant and Young Child Feeding
AWD	Acute Watery Diarrhoea	Kcal	Kilocalorie(s)
AWG	Average Weight Gain	Kg	Kilogram(s)
BSFP	Blanket Supplementary Feeding Programme	L	Litre(s)
CBNC	Community Based Newborn Care	LOS	Length of Stay
cm	Centimetre(s)	LRTI	Lower Respiratory Tract Infection
CSAS	Centric Systematic Area Sampling	µg	Microgram(s)
CTC	Cholera Treatment Centre	MAM	Moderate Acute Malnutrition
dL	Deci-litre(s)	mg	Milligram(s)
EPI	Expanded Programme on Immunisation	ml	Millilitre(s)
ETAT	Emergency Triage Assessment and Treatment	mm	Millimetre(s)
FBF	Fortified Blended Foods	FMOH	Federal Ministry of Health
g	Gram(s)	MUAC	Mid-Upper Arm Circumference
GAM	Global Acute Malnutrition	NGT	Naso-Gastric Tube
GFD	General Food Distribution	NNP	National Nutrition Programme
GMP	Growth Monitoring Promotion	ORS	Oral Rehydration Solution
Hb	Haemoglobin	OTP	Outpatient Therapeutic Programme
HC	Health Centre	PFSA	Pharmaceutical Fund and Supply Agency
HDA	Health Development Army	PHCU	Primary Health Care Unit
HDG	Health Development Group	PLHIV	Person or People Living with HIV
HEW	Health Extension Worker	PLW	Pregnant and Lactating Women
HIV	Human Immunodeficiency Virus	PSNP	Productive Safety Net Programme
HMIS	Health Management Information System	ReSoMal	Rehydration Solution for Malnutrition
HP	Health Post	RHB	Regional Health Bureau
HSTP	Health Sector Transformation Plan	RUSF	Ready-to-Use Supplementary Food

RUTF	Ready-to-Use Therapeutic Food
S3M	Simple Spatial Survey Method
SAM	Severe Acute Malnutrition
SBCC	Social Behaviour Change Communication
SC	Stabilisation Centre
SFP	Supplementary Feeding Programme
SLEAC	Simplified Lot quality assurance sampling Evaluation of Access and Coverage
SQUEAC	Semi-Quantitative Evaluation of Access and Coverage
TSFP	Targeted Supplementary Feeding Programme
TB	Tuberculosis
UNICEF	United Nations International Children's Emergency Fund
WASH	Water Sanitation and Hygiene
WFP	World Food Programme
WFA	Weight-For-Age
WFL	Weight-For-Length
WFH	Weight-For-Height
WHO	World Health Organisation
ZHD	Zonal Health Department



Definition of Terms

Term	Definition
Acute Malnutrition	Acute malnutrition is a form of undernutrition. It is caused by a decrease in food consumption and/or illness resulting in bilateral pitting oedema and/or sudden weight loss. It is defined by the presence of bilateral pitting oedema or wasting [low Mid-Upper Arm Circumference (MUAC) or low Weight-For-Height (WFH)/Weight-For-Length (WFL)].
Anthropometry	Anthropometry is the study and technique of human body measurement. It is used to measure and monitor the nutritional status of an individual or population group.
Appetite Test	Appetite test is the decisive criteria for participation in Outpatient Therapeutic Programme (OTP). The test is done at admission and at all OTP follow-on visits to ensure that the child 6–59 months or older can eat Ready-to-Use Therapeutic Food (RUTF). If the patient has no appetite, s/he must be referred to the Stabilisation Centre (SC).
Blanket Supplementary Feeding Programme (BSFP)	BSFP is an intervention that aims to prevent acute malnutrition among vulnerable groups. A supplementary food ration is provided to everyone in an identified vulnerable group for a defined period. This might be all children 6–24 or 6–59 months old and/or all Pregnant and Lactating Women (PLW), regardless of their nutritional status.
Bilateral Pitting Oedema	Bilateral pitting oedema, also known as kwashiorkor or oedematous malnutrition, is a sign of Severe Acute Malnutrition (SAM). It is defined by bilateral pitting oedema of the feet and verified when thumb pressure applied on top of both feet for three seconds leaves a pit (indentation) in the feet after the thumb is lifted. It is an abnormal infiltration and excess accumulation of serous fluid in connective tissue or in a serous cavity. Grades of bilateral pitting oedema are + (mild), ++ (moderate) and +++ (severe).
Community-Based Management of Acute Malnutrition (CMAM)	CMAM refers to the community-based management of acute malnutrition through: 1) SC for the management of SAM with medical complications; 2) OTP for the management of SAM without medical complications; 3) community outreach; and 4) services or programmes for the management of Moderate Acute Malnutrition (MAM).

Community Outreach

Community outreach for CMAM is a continuous process that includes the following activities: community assessment, formulating a community outreach strategy, developing and disseminating messages and materials, training on community service providers, community mobilisation, active case-finding and referral, home visits and follow-up, and linking with other community programmes and initiatives.

F75

Formula 75 (75 kcal/100 ml) is a therapeutic milk recommended by WHO for the stabilisation phase of patients with SAM and medical complications.

F100

Formula 100 (100 kcal/100ml) is a therapeutic milk recommended by WHO for nutrition rehabilitation of children with SAM after the stabilisation phase in the SC. F100 has a similar nutrient composition to RUTF.

F100 Diluted

Formula 100 Diluted (130 kcal/100ml) is a therapeutic milk recommended by WHO for the stabilisation and nutrition rehabilitation of infants 0-6 months with SAM and without bilateral pitting oedema in the SC.

Fortified Blended Foods (FBF)

FBF are a mixture of cereals and other ingredients such as soya beans or pulses that have been milled, blended, pre-cooked by extrusion or roasting, and fortified with a pre-mix of a sufficient amount and range of vitamins and minerals. Super Cereal Plus is an example of FBF.

GAM

GAM is a population-level indicator referring to overall acute malnutrition defined by the presence of bilateral pitting oedema or wasting defined by WFH < -2 z-score (WHO standards). GAM is a sum of SAM and MAM (GAM = SAM + MAM).

Height-for-Age (HFA) Index

The HFA index is used to assess stunting. It shows how a child's height compares to the height of a child of the same age and sex in the WHO standards. This index reflects a child's past nutritional status.

Moderate Acute Malnutrition (MAM)

MAM, or moderate wasting, is defined by a MUAC \geq 11.5 cm and < 12.5 cm or a WFH \geq -3 z-score and < -2 z-score (WHO standards) in children 6-59 months old.

MAM can also be used as a population-level indicator defined by WFH \geq -3 z-score and < -2 z-score (WHO standards).

MUAC Indicator

Low MUAC is an indicator for wasting. In a child that is 6–59 months of age, MUAC < 11.5 cm indicates severe wasting, or SAM. MUAC \geq 11.5 cm and < 12.5 cm indicates moderate wasting, or MAM.

Outpatient Therapeutic Care (OTP) for Management of SAM Without Medical Complications

OTP is a service treating patients with SAM without medical complications through the provision of routine medical treatment and nutrition rehabilitation with RUTF. Patients attend OTP every week until the discharge criteria are reached.

Ready-to-Use
Supplementary Food
(RUSF)

RUSF is an energy-dense, mineral- and vitamin-enriched formulation specifically designed to treat MAM. The composition of RUSF is different to that of RUTF which is designed to treat SAM. RUSF is soft and can be consumed easily by children from the age of 6 months without adding water. It is not water-based, meaning that bacteria cannot grow in it and that it can be used safely at home without refrigeration and in areas where hygiene conditions are not optimal. Unlike fortified blended foods (FBF), RUSF does not require preparation before consumption. Plumpy'sup® is an example of a known lipid-based RUSF.

Ready-to-Use
Therapeutic Food (RUTF)

RUTF is an energy-dense, mineral- and vitamin-enriched formulation specifically designed to treat SAM. RUTF has a similar nutrient composition to F100. RUTF is soft and can be consumed easily by children from the age of 6 months without adding water. Unlike F100, RUTF is not water-based, meaning that bacteria cannot grow in it and it can be used safely at home without refrigeration and in areas where hygiene conditions are not optimal. It does not require preparation before consumption. Plumpy'nut® is an example of a known lipid-based RUTF.

Severe Acute
Malnutrition (SAM)

SAM is defined by the presence of bilateral pitting oedema or severe wasting (MUAC < 11.5 cm or a WFH < -3 z-score [WHO standards]) in children 6-59 months old.

A patient with SAM is highly vulnerable and has a high mortality risk.

SAM can also be used as a population-based indicator defined by the presence of bilateral pitting oedema or severe wasting (WFH < -3 z-score [WHO standards]).

Severe wasting is a sign of SAM. It is defined by a MUAC < 11.5 cm or a WFH < -3 z-score [WHO standards]) in children 6-59 months.

Severe Wasting

Severe wasting is also called marasmus. The patient with severe wasting has lost fat and muscle and appears very thin (e.g., signs of "old man face" or "baggy pants" [folds of skin over the buttocks]).

Severe Wasting with
Bilateral Pitting Oedema

Severe wasting with bilateral pitting oedema is the simultaneous condition of severe wasting (MUAC < 11.5 cm or WFH < -3 in children 6-59 months) and bilateral pitting oedema (of any grade +, ++ or +++).

Sphere Project or
Sphere Standards

The Sphere Project Humanitarian Charter and Minimum Standards in Disaster Response is a voluntary effort to improve the quality of assistance provided to people affected by disaster and to enhance the accountability of the humanitarian agencies in disaster response. Sphere has established Minimum Standards in Disaster Response (often referred to as Sphere Standards) and indicators to describe the level of disaster assistance needed. Visit www.sphereproject.org for more information.

Stabilisation Centre (SC)

SC is a service treating patients with SAM with medical complications until their medical conditions are stabilised and complications are resolved (usually four to seven days). Treatment then continues in OTP until the discharge criteria are reached. SC for SAM with medical complications is provided in a Health Centre or Hospital with 24-hour care capacity.

Stunting

Stunting, or chronic undernutrition, is a form of undernutrition and presents as low height-for-age (HFA). It is defined by an HFA z-score below two SDs of the median (WHO standards). Stunting is a result of prolonged or repeated episodes of undernutrition which can start before birth.

Targeted Supplementary Feeding Programme (TSFP)

TSFP is an intervention that aims to treat patients with MAM, and prevents deterioration of the condition to SAM. The supplementary food ration is targeted to individuals with MAM in specific vulnerable groups such as children age 6-59 months and malnourished pregnant women and lactating women with infants under 6 months of age.

Undernutrition

Undernutrition is a consequence of a deficiency in nutrient intake and/or absorption in the body. The different forms of undernutrition that can appear isolated or in combination are acute malnutrition (bilateral pitting oedema and/or wasting), stunting, underweight (includes elements of wasting and stunting), and micronutrient deficiencies.

Underweight

Underweight is a composite form of undernutrition including elements of stunting and wasting and is defined by a weight-for-age (WFA) < -2 z-score (WHO standards).

Wasting

Wasting is a form of acute malnutrition. It is defined by a MUAC < 12.5 cm or a WFH < -2 z-score (WHO standards) in children 6-59 months.

WFA Index

The WFA index is used to assess underweight. It shows how a child's weight compares to the weight of a child of the same age and sex in the WHO standards. The index reflects a child's combined current and past nutritional status.

WFH/WFL Index

The WFH/WFL index is used to assess wasting. It shows how a child's weight compares to the weight of a child of the same height/length and sex in the WHO standards. The index reflects a child's current nutritional status.

1 Introduction

Ethiopia is prone to recurrent droughts that impact food security and water availability, and contribute to high rates of malnutrition in the country. Over the past decade, the Government has applied a two-pronged approach to address malnutrition. The first aspect of the approach focuses on increasing access and availability to food through improved economic growth, better agricultural production systems along with promotion of good nutrition practices and prevention of malnutrition. The second aspect aims to strengthen early warning systems and timely emergency response, including wide-scale delivery of services for the management of acute malnutrition.

This *National Guideline for the Management of Acute Malnutrition in Ethiopia* aims to improve access to quality services for Severe Acute Malnutrition (SAM) and Moderate Acute Malnutrition (MAM). It incorporates the latest international and national evidence on acute malnutrition. The guideline replaces the Protocol for the Management of SAM (2007), and the Guideline for the Management of MAM (2012). It also partially replaces content provided in the Guideline for HIV/AIDS and Nutrition (2010).

Purpose and Scope of the Guideline

The purpose of this guideline is to provide harmonised practical direction to service providers, policy makers, programmers, and stakeholders on the current standards and recommendations on acute malnutrition. The guideline also aims to:

- ◆ Prevent deterioration of acute malnutrition through early identification, referral, follow-up, and referring patients to other health and nutrition interventions.
- ◆ Treat acute malnutrition to reduce associated morbidity and mortality.

The guideline is aligned to the National Nutrition Programme (NNP) II 2016-2020 and the Health Sector Transformation Plan (HSTP) 2015/16 - 2019/20. It targets children 0-59 months, Pregnant and Lactating Women (PLW), older people (≥ 60 years), and other vulnerable groups including; HIV/AIDS, tuberculosis (TB), and disabilities. The guideline complements other health and nutrition interventions and should be used alongside the following documents:

-
1. *Comprehensive Integrated Nutrition Services, 2016*
 2. *National Guideline on Adolescent, Maternal, Infant and Young Child Nutrition, 2016*
 3. *National Implementation Guideline for Integrated Community Case Management of Childhood Illnesses and Newborn Care, 2017*
 4. *National Guidelines for Comprehensive HIV Prevention, Treatment and Care, 2018/19*
 5. *National Guidelines for Management of Tuberculosis (TB), Drug Resistant-TB and Leprosy in Ethiopia, 2017*
 6. *National Malaria Guidelines, 2018*
 7. *Standard Treatment Guidelines for General Hospitals, 2014*
 8. *Standard Treatment Guidelines for Primary Hospitals, 2014*
 9. *Standard Treatment Guidelines for Health Centres, 2014*

Use of the Guideline

The guideline should be used by facility- and community-based service providers. It can also be used by policymakers, programme advisors, coordinators, and managers working at national and sub-national levels to design, implement, monitor, and evaluate nutrition programmes.

The National Guideline for the Management of Acute Malnutrition in Ethiopia comes with a set of implementation tools including **quick reference guides**, **monitoring and reporting tools**, and **training materials**.

What is New in The Guideline

The table below summarises changes made to the revised guideline in comparison to the Protocol for the Management of SAM (2007), and the Guideline for the Management of MAM (2012). The changes reflect most recent evidence and best practices on the management of acute malnutrition.

1 Admission and Discharge Criteria for Children 6-59 Months	
Old	SAM admission was based on MUAC <11.0 cm and any degree of bilateral pitting oedema. Children receiving SAM treatment were discharged when they reach a WFH percentage of the median of 85% for two consecutive visits.
New	Admission for SAM treatment is now based on MUAC <11.5 cm or any degree of bilateral pitting oedema or WFH <-3 z-score; for MAM treatment a MUAC of ≥11.5 to <12.5 cm or WFH ≥-3 to <-2 z-score is used. Children receiving SAM or MAM treatment should be discharged when they reach a MUAC ≥ 12.5 cm or WFH ≥ -2 z-scores and have no bilateral pitting oedema for two consecutive visits.
Old	Used the NCHS reference standards WFH percentage of the median to determine admission for treatment.
New	If WFH is used, only use the WFH z-score of the WHO growth standards.
Old	WFH percentage of the median was used to determine discharge regardless of the criteria used to admit the patient for the management of SAM.
New	<ul style="list-style-type: none"> ◆ The anthropometric indicator that is used to confirm SAM should also be used to assess whether a child has reached nutritional recovery, i.e., if MUAC is used to identify that a child has SAM, then MUAC should be used to assess and confirm nutritional recovery and discharge. Similarly, if WFH is used to identify that a child has SAM, and then WFH should be used to assess and confirm nutritional recovery and discharge. ◆ Children admitted with only bilateral pitting oedema should be discharged based on either MUAC or WFH. ◆ Percentage weight gain should not be used as a discharge criterion.
2 Use of Antibiotics in Children 0-59 Months	
Old	Cotrimoxazole or amoxicillin were used as routine medication for SAM without medical complications.
New	Only amoxicillin is used as a routine medication for all patients with SAM without medical complications. If a patient is HIV positive and receiving cotrimoxazole according to the National Guidelines for Comprehensive HIV Treatment and Care, then amoxicillin is given in addition as a routine SAM medication.
Vitamin A Supplementation for Children 0-59 Months	
Old	A high dose of vitamin A was routinely given to all children with SAM regardless of the therapeutic food they were receiving.
New	<ul style="list-style-type: none"> ◆ A high dose of vitamin A is not given to patients if they are receiving therapeutic foods that comply with WHO specifications. A high dose of vitamin A is given on admission only if the therapeutic foods provided are not fortified as recommended in the WHO specifications and vitamin A is not part of other daily supplements. ◆ If the patient with SAM has eye signs of vitamin A deficiency or recently had measles (within the past 3 months), a high dose of vitamin A (50,000 IU, 100,000 IU or 200,000 IU depending on age) is given on day 1, 2, and 15, regardless of the therapeutic food they are receiving.

3 Folic Acid Supplementation for Children 0-59 Months	
Old	On the day of admission, a single dose of folic acid (5mg) was given to children with SAM and clinical signs of anaemia.
New	Folic acid should not be given routinely to SAM children receiving therapeutic foods (F-75, F-100 or RUTF) that comply with WHO specifications, as they contain enough amounts of folic acid.
4 Therapeutic Feeding Approach for SAM with Medical Complications During Transition Period	
Old	Either F-100 or RUTF was recommended for use during transition.
New	During the transition period, RUTF is introduced alongside F-75. The child drinks water freely. If the child does not consume the prescribed amount of RUTF, F-75 is used to top-up the feed. The amount of RUTF is increased over 2–3 days until the child consumes the full required amount of RUTF.
5 Fluid Management of with SAM with Medical Complications	
Old	<ul style="list-style-type: none"> ◆ Children 0-59 months with SAM who presented with dehydration but not in shock were rehydrated slowly, either orally or by Naso-Gastric Tube (NGT) using Rehydration Solution for Malnutrition (ReSoMal). ◆ Oral Rehydration Solution (ORS) was not used for oral or nasogastric rehydration of children with SAM. ◆ For patients with SAM and signs of shock or severe dehydration who could not be rehydrated orally or by NGT, IV fluids of half-strength Darrow’s solution with 5% glucose or Ringer’s lactate solution with 5% glucose were used for treatment. If not available, 0.45% saline with 5% glucose were used.
	<ul style="list-style-type: none"> ◆ Children 0-59 months with SAM who present with dehydration but not in shock should be rehydrated slowly, either orally or by NGT using ReSoMal (5-10ml/kg/h up to a maximum of 12 hours). ◆ ORS should not be used for oral or nasogastric rehydration in children with SAM with dehydration. ◆ Patients with SAM and signs of shock or severe dehydration and who cannot be rehydrated orally or by NGT should be treated with IV fluids of Ringer’s lactate solution with 5% dextrose. If not available, 0.45% saline and 5% dextrose should be used. ◆ Low osmolarity ORS is recommended instead of ReSoMal if a patient is diagnosed with profuse watery diarrhoea or acute watery diarrhoea (AWD). ◆ Guidance has been provided on the management of AWD in patients with SAM (see Annex 2 for details).
	New

6 Management of Acute Malnutrition in the Context of HIV/AIDS

Old	<ul style="list-style-type: none"> ◆ Same treatment of malnutrition was recommended for all patients with SAM irrespective of their HIV status. ◆ The treatment of malnutrition was started at least one week before the introduction of Anti-Retroviral Therapy (ART) to diminish the risk of serious side effects from the drugs. ◆ Children with HIV were given co-trimoxazole preventive therapy. In addition, amoxicillin was also given.
New	<ul style="list-style-type: none"> ◆ All patients failing to respond to SAM or MAM treatment should be offered HIV counselling and testing. ◆ Refer HIV-positive patients with SAM and MAM for treatment according to National Guidelines for Comprehensive HIV Prevention, Treatment and Care. ◆ HIV-positive children with SAM or MAM should be given the same antiretroviral therapy (ART) treatment regimen, in the same doses, as patients with HIV who do not have SAM or MAM. ◆ Children with SAM or MAM who are HIV-positive should be managed using the same specialised nutritious foods (i.e., F-75, F-100, RUTF, RUSE, or FBF) as those with SAM or MAM who are HIV-negative. ◆ Children with SAM and MAM who are HIV-positive should follow the same guidance for micronutrient supplementation including Vitamin A, iron, folic acid and zinc for diarrhoea management as those with SAM and MAM who are HIV-negative. ◆ Children with SAM who are HIV-positive and qualify for ART should be started on treatment as soon as possible after stabilisation. ◆ HIV-positive children with SAM who are started on ART treatment should be monitored closely 6–8 weeks following initiation of ART. ◆ HIV-positive children with SAM in whom persistent diarrhoea does not resolve with standard management should be investigated to exclude carbohydrate intolerance and infective causes, which may require different management, such as modification of fluid and feed intake, or antibiotics.

7 Management of Infants 0-6 Months of Age

Old	<p>Infants 0-6 months with a WFL of < 70% of the percentage of the median and visible signs of wasting or bilateral pitting oedema were referred for the management of SAM in the SC.</p>
New	<p>Infants 0-6 months of age should be managed in SC if they meet the following criteria:</p> <ul style="list-style-type: none"> ◆ Any grade of bilateral pitting oedema (+, ++ or +++) ◆ WFL < -3 z-score ◆ Any serious clinical conditions or medical complications as outlined for children 6-59 months. ◆ Recent weight loss or failure to gain weight. ◆ Ineffective feeding (attachment, positioning and suckling) directly observed for 15-20 min, ideally in a supervised, separated area. ◆ Any medical or social issue needing more detailed assessment or intensive support (e.g., disability, depression of caregiver, or other adverse social circumstances).

8 Management of MAM in Children 6-59 Months and PLW

Old Targeted Supplementary Feeding Programme (TSFP) was implemented at the community level. Services were delivered by Non-Governmental Organisations (NGOs) and not integrated with routine health services.

New

- ◆ TSFP will be integrated to routine health services in priority areas identified by the FMOH.
- ◆ TSFP should be integrated with the treatment of SAM, particularly in OTP, to ensure a continuum of care for acute malnutrition.

9 Management of Acute Malnutrition in Older People (≥ 60 years)

Old No guidance was provided on the management of acute malnutrition in the older people.

New This guideline provides guidance on the management of acute malnutrition in older people. Care and treatment follow the same protocols as that of children 6-59 months, below is the admission criteria for older people:

- ◆ MUAC < 18.5 cm admit for management of SAM in Outpatient Therapeutic Programme (OTP).
- ◆ MUAC ≥ 18.5 to 21.0 cm admit for management of MAM in TSFP.

10 Supply Management

Old No guidance was provided on the integrated supply management of SAM and MAM commodities.

New This guideline provides guidance on the supply management of SAM and MAM commodities and facilitates future integration into the Integrated Pharmaceutical Logistics Systems (IPLS). Guidance provided includes requesting and receiving supplies, stock management, and reporting of supplies.

2 Integrating Acute Malnutrition Services into the Health System

This chapter highlights the core aspects of the care and treatment for SAM and MAM, and integration into the routine health system. The chapter also summarises the roles and responsibilities of the stakeholders engaged with the management of acute malnutrition in the Ethiopian context.

2.1 Acute Malnutrition in the Context of Ethiopia

In line with the NNP II 2016-2020 and the HSTP 2015/16-2019/20, services for the management of SAM and MAM should be delivered through the health system including; Health Posts (HP), Health Centres (HC), Woreda, Zonal, Regional and referral Hospitals.

Community and Household Level

Good community outreach is critical in ensuring early identification of SAM and MAM cases. Community outreach also aims to empower communities and families to understand the causes of malnutrition, and prevent and manage acute malnutrition in their communities.

For maximum impact, community outreach activities should be integrated with other community-based health and nutrition interventions. These include Integrated Management of Newborn and Childhood illness (IMNCI), Integrated Community Case Management and Community Based Newborn Care (ICCM-CBNC), child health campaigns, Infant and Young Child Feeding (IYCF) and Growth Monitoring Promotion (GMP). Community outreach activities should also be linked to multi-sectoral nutrition programmes, and initiatives to eradicate poverty and improve food security; Water, Sanitation and Hygiene (WASH), and social protection.

Health Extension Workers (HEWs) collaborate with the community-based structures such as health committees and engage with the Health Development Army (HDA)/Health Development Group (HDG) to screen and refer cases to the appropriate service for treatment. See Figure 1, the flow chart on the management of acute malnutrition.

Health Post Level

The HPs provide primary health care services such as disease prevention and control, hygiene and environmental sanitation, family health services, and health education and communication. The service provider at the HP works closely with the network of the HDA/HDG to facilitate the management of SAM and MAM. The service provider diagnoses acute malnutrition and provides Outpatient Therapeutic Programme (OTP) and Targeted Supplementary Feeding Programme (TSFP) services. They also determine patients who have medical complications and refer them to the Stabilisation Centre (SC)


Health Centre Level

The HC provides 24-hour SC for the management of SAM with medical complications. The HC also manages and provides technical support to a cluster of one to five HPs within its vicinity.

Hospital Level

Woreda, Zonal, Regional and referral Hospitals provide higher level referral points where further care can be provided to patients with SAM and more complex medical complications. The referral Hospitals have the facilities and expertise to manage situations that may require administration of oxygen, blood transfusion, and other critical care.

Figure 1. Flow Chart on the Management of Acute Malnutrition



* **Medical complications include:** Poor appetite, intractable vomiting, convulsions, lethargy, not alert, unconsciousness, high fever (axillary temperature >39 °C), lower respiratory tract infection, dehydration, persistent diarrhoea, severe anaemia, hypoglycaemia, hypothermia (axillary temp. < 35 °C), severe skin lesions, eye signs of vitamin A deficiency.

Table 1. Assessing and Classifying Acute Malnutrition in Infants 0-6 Months

Ask	Look and Feel	Signs	Classify	Treat
<p>Ask for signs of medical complications:</p> <p>Is the patient:</p> <ul style="list-style-type: none"> ■ Too weak to suckle effectively? ■ Vomiting everything? <p>Does the patient have:</p> <ul style="list-style-type: none"> ■ Recent weight loss or failure to gain weight? <p>Does the patient have:</p> <ul style="list-style-type: none"> ■ Blood in stool? ■ Diarrhoea > 14days? ■ Cough > 14 days, or had contact with TB patients? ■ Bleeding tendencies? ■ History of recent sunken eyes? ■ Convulsion(s) (more than one or prolonged for > 15 min)? 	<p>1. Check for presence of bilateral pitting oedema</p> <ul style="list-style-type: none"> ■ Does the infant have bilateral pitting oedema? <p>2. Check the WFL</p> <ul style="list-style-type: none"> ■ Is the infant's WFL < -3 z-score? <p>3. Check for medical complications:</p> <ul style="list-style-type: none"> ■ Poor appetite ■ Intractable vomiting ■ Convulsions ■ Lethargy, not alert ■ Unconsciousness ■ High fever (axillary temp. >39 °C) ■ Lower respiratory tract infection ■ Dehydration ■ Persistent diarrhoea ■ Severe anaemia ■ Hypoglycaemia, hypothermia (axillary temp. <35 °C) ■ Severe skin lesions ■ Eye signs of vitamin A deficiency 	<ul style="list-style-type: none"> ■ Any grade of bilateral pitting oedema (+, ++ or +++) OR ■ WFL < -3 z score OR: ■ Recent weight loss or failure to gain weight. OR: ■ Ineffective feeding (attachment, positioning and suckling) directly observed for 15-20, minutes, ideally in supervised separate area. OR: ■ Presence of any of the following medical complications: <ul style="list-style-type: none"> ◆ Poor appetite ◆ Intractable vomiting ◆ Convulsions ◆ Lethargy, not alert ◆ Unconsciousness ◆ High fever ◆ Lower respiratory tract infection ◆ Dehydration ◆ Persistent diarrhoea ◆ Severe anaemia ◆ Hypoglycaemia ◆ Hypothermia ◆ Severe skin lesions ◆ Eye signs of vitamin A deficiency 	<p>SAM with and without medical complications.</p>	<p>Refer/Admit in SC.</p>

	<p>4. Check for any other medical or social issues?</p> <ul style="list-style-type: none"> Does the infant have medical or family issues that need more detailed assessment or support? <p>5. Check the mother's MUAC.</p> <ul style="list-style-type: none"> Is the mother's MUAC <23.0cm? 	<p>OR:</p> <ul style="list-style-type: none"> Any medical or social issue needing more detailed assessment or intensive support (e.g. disability, depression of caregiver, or other adverse social circumstances). 		<p>Assess the infant's feeding and counsel the mother or caregiver on appropriate IYCF practices. Emphasize on establishing effective exclusive breastfeeding.</p> <ul style="list-style-type: none"> If feeding problems, follow up in 5 days. If no feeding problem, follow up in 30 days. Admit the mother to TSFP. Refer the mother for Productive Safety Net Programme (PSNP).
	<ul style="list-style-type: none"> WFH ≥ -3 to < -2 Z scores No bilateral pitting oedema AND No medical complications Clinically well and alert Mother has a MUAC of < 23.0 cm 	MAM		<p>Congratulate and counsel the mother on appropriate IYCF practices.</p>
	<ul style="list-style-type: none"> WFH ≥ -2 Z score AND No bilateral pitting oedema 	No acute malnutrition.		

Table 2. Assessing and Classifying Acute Malnutrition in Children 6-59 Months

Ask	Look and Feel	Signs	Classify	Treat
<p>Ask for signs of medical complications:</p> <p>Is the patient:</p> <ul style="list-style-type: none"> ■ Too weak to suckle effectively? ■ Vomiting everything? <p>Does the patient have:</p> <ul style="list-style-type: none"> ■ Blood in stool? ■ Diarrhoea > 14days? ■ Cough > 14 days, or had contact with TB patients? ■ Bleeding tendencies? ■ History of recent sunken eyes? ■ Convulsion(s) (more than one or prolonged for > 15 min)? 	<p>1. Check for presence of bilateral pitting oedema</p> <ul style="list-style-type: none"> ■ Does the infant have bilateral pitting oedema? ■ If yes, is it +++ (generalised involving upper arms and face)? <p>2. Check MUAC</p> <ul style="list-style-type: none"> ■ Is the child's MUAC < 11.5cm? <p>3. Check the WFL</p> <ul style="list-style-type: none"> ■ Is the infant's WFL < -3 z-score? <p>4. Check for medical complications:</p> <ul style="list-style-type: none"> ■ Poor appetite ■ Intractable vomiting ■ Convulsions ■ Lethargy, not alert ■ Unconsciousness ■ High fever (>39 °C) ■ Lower respiratory tract infection ■ Dehydration 	<ul style="list-style-type: none"> ■ Any grade of bilateral pitting oedema (+, ++ or +++) OR ■ WFL < -3 z score OR: ■ Recent weight loss or failure to gain weight. OR: ■ Ineffective feeding (attachment, positioning and suckling) directly observed for 15-20, minutes, ideally in supervised separate area. OR: ■ Presence of any of the following medical complications: <ul style="list-style-type: none"> ◆ Poor appetite ◆ Intractable vomiting ◆ Convulsions ◆ Lethargy, not alert ◆ Unconsciousness ◆ High fever ◆ Lower respiratory tract infection ◆ Dehydration ◆ Persistent diarrhoea ◆ Severe anaemia 	SAM with and without medical complications.	Refer/Admit in SC.

<ul style="list-style-type: none"> ■ Persistent diarrhoea ■ Severe anaemia ■ Hypoglycaemia, hypothermia (axillary temp. <35 °C) ■ Severe skin lesions ■ Eye signs of vitamin A deficiency 	<ul style="list-style-type: none"> ◆ Hypoglycaemia ◆ Hypothermia ◆ Severe skin lesions ◆ Eye signs of vitamin A deficiency 	<p>SAM without medical complications</p>	<p>Admit in OTP. (See section 4.2 for the treatment and care plan)</p>
	<ul style="list-style-type: none"> ■ Bilateral pitting oedema + or ++ OR ■ MUAC <11.5 cm OR ■ WFH <-3 z-score AND ■ Appetite test passed ■ No medical complications ■ Clinically well and alert 	<p>MAM</p>	<p>Admit in TSFP and counsel on appropriate IYCF practices (See chapter 5.0 for the treatment and care plan)</p>
	<ul style="list-style-type: none"> ■ MUAC ≥11.5 to <12.5 cm ■ WFH ≥ -3 to <-2 z scores ■ No bilateral pitting oedema AND ■ No medical complications ■ Clinically well and alert 	<p>No acute malnutrition.</p>	<p>Congratulate and counsel the mother on appropriate IYCF practices.</p>
	<ul style="list-style-type: none"> ■ MUAC ≥ 12.5 cm ■ WFH ≥ -2 z score AND ■ No bilateral pitting oedema 		

2.2 Roles and Responsibilities of Stakeholders

Described here are the roles and responsibilities of the various stakeholders involved in the management of acute malnutrition within the Ethiopian health system.

2.2.1 The Health System

The Federal Ministry of Health (FMOH)

- ◆ Provides national guidance on the management of acute malnutrition, in line with the NNP II 2016-2021.
- ◆ Ensures that adequate numbers of skilled health professionals are assigned to health facilities.
- ◆ Ensures programme quality through monitoring and supervision.
- ◆ Coordinates national activities including orientations, trainings, and consultation workshops.
- ◆ Mobilises resources for the implementation of SAM and MAM activities.
- ◆ Ensures continuous supply of essential commodities to all Regional Health Bureaus (RHBs), [see Annex 13 and 14](#) for the list of essential SAM and MAM supplies.
- ◆ When necessary, reviews policy and implementation guidelines.

The Regional Health Bureaus (RHBs) and Zonal Health Department (ZHD)

- ◆ Provides programme implementation guidance to the Zonal Health Department (ZHD) and Woreda health offices.
- ◆ Coordinates and supports implementation of training at HC and HP level.
- ◆ Conducts monitoring and supervision to the HCs, and HPs at the Woreda level.
- ◆ Mobilises resources for trainings at the Regional, Zonal and Woreda level.
- ◆ Ensures continuous supply of essential commodities to all Zones and Woredas, [see Annex 13 and 14](#) for the list of essential SAM and MAM supplies.
- ◆ Ensures quality and timely reporting of activities within the Region and Zone.

The Woreda Health Office

- ◆ Coordinates and supports training of health care providers and HEWs.
- ◆ Ensures continuous supply of essential commodities to all health facilities, [see Annex 13 and 14](#) for the list of essential SAM and MAM supplies.

-
- ◆ Strengthens the referral and communication system between HCs and HPs.
 - ◆ Ensures that the HC service providers conduct regular supportive supervision to the HPs.
 - ◆ Conducts supportive supervision and regular review meetings with HCs and HPs.
 - ◆ Ensures timely reporting of activities to the ZHD.

The Primary Health Care Unit (PHU) and Referral Health Centre (HC)

- ◆ Coordinates and participates in training of health care providers and HEWs.
- ◆ Builds the capacity of HEWs on the management of acute malnutrition.
- ◆ Ensures a continuous supply of specialised nutritious foods to all HPs within the Kebele catchment area.
- ◆ Conducts supportive supervision to the HPs.
- ◆ Ensures that IMNCI/ICCM-CBNC activities are implemented and coordinated within the Kebele catchment areas.
- ◆ Submits accurate and timely Monthly Statistics Reports for the Management of Acute Malnutrition.

The Health Post (HP) Level

- ◆ Ensures quality implementation of SAM and MAM activities.
- ◆ Ensures that SAM and MAM services are delivered routinely through the health care system.
- ◆ Ensures continuous availability and proper utilization of the essential supplies for SAM and MAM, [see Annex 13 and 14](#) for the list of essential SAM and MAM supplies.
- ◆ Ensures that IMNCI/ICCM-CBNC activities are implemented at the HP.
- ◆ Prepares and submits accurate and timely Monthly Statistics Reports for the Management of Acute Malnutrition.
- ◆ Engages with and builds the capacity of the HDA/HDG on how to conduct community mobilisation, referral, follow-up, counselling, and support of patients with acute malnutrition.
- ◆ Collaborates with the HDA/HDG to conduct home visits, follow-up and referrals of SAM and MAM patients.
- ◆ Ensures that PLW and families with SAM and MAM patients are enrolled to the Productive Safety Net Programme (PSNP) and that they understand the conditions for the support they receive.

The Health Development Army/Health Development Group (HDA/HDG)

- ◆ Conducts health promotion and counselling at the community level.
- ◆ Mobilises communities and refers caregivers to the HEW for screening of acute malnutrition.
- ◆ Counsels and supports caregivers ensuring that the referred cases report to the HP.
- ◆ Counsels and supports the caregiver with treatment compliance on the consumption of specialised nutritious foods and routine medications.

2.2.2 The United Nations (UN) and Donor Agencies

- ◆ Advocates for the management of acute malnutrition, and children's rights to food and nutrition.
- ◆ Contributes to mobilisation of resources for implementation of prioritised activities.
- ◆ Provides funds to implement the National Guideline for the Management of Acute Malnutrition.
- ◆ Provides technical support to FMOH and RHB to facilitate the management of acute malnutrition in an integrated manner.
- ◆ Supports the FMOH and RHB to implement their respective roles and responsibilities.

2.2.3 Non-governmental Organisations (NGOs) and Community-Based Organisations (CBOs)

- ◆ Advocates for the management of acute malnutrition, and children's rights to food and nutrition.
- ◆ Provides technical support to the ZHD, Woreda health office, HCs, HPs, and communities to facilitate the management of acute malnutrition in an integrated manner.
- ◆ Supports ZHD, Woreda health office, HCs and HPs to implement their respective roles and responsibilities.
- ◆ Where possible, provides direct support to families and communities, especially those under difficult circumstances.

3 Community Outreach

This chapter highlights why community involvement and empowerment are critical to improving access and uptake of acute malnutrition services. The chapter also describes the core components of community outreach activities and those who should be involved in service delivery.

3.1 Planning Community Outreach

3.1.1. Community Assessments

In the case of sudden increase or decrease in the SAM and MAM case load, the HEWs, under the guidance of the Woreda health office and PHCU nutrition focal person should conduct a community assessment to understand if acute malnutrition services are meeting the demand of the population at the Kebele level. The community assessment should:

- ◆ Engage with the HDA/HDG with 30 households organised into groups of one to five women to:
 - ◆ Share data on acute malnutrition in the area.
 - ◆ Explain any increases or decreases in the levels of acute malnutrition.
 - ◆ Discuss the causes and possible solutions to addressing acute malnutrition in the community.
- ◆ Engage with the health committees, agriculture extension workers, teachers, and other community workers to identify opportunities for collaborating in addressing acute malnutrition in the area.
- ◆ Identify available initiatives and non-governmental organisations supporting health, nutrition and food security activities in the area.
- ◆ Identify barriers and enablers to nutrition and health including attitudes and health-seeking behaviours.

3.1.2. Formulation of a Community Outreach Strategy

- ◆ Identify the best mechanisms to improve community outreach activities.
- ◆ Organise meetings with the community and religious leaders to ask for their continued support in acute malnutrition services.
- ◆ Develop a plan and define the roles and responsibilities for the people who will be involved in community outreach and mobilisation activities.

3.1.3. Develop Messages and Materials

- ◆ Develop simple and clear messages that are easy to understand. Graphic images or pictures should be used to explain complex concepts.
- ◆ Disseminate acute malnutrition messages using the HDA/HDG structure, and other health and nutrition information communication channels.
- ◆ Methods of disseminating information may include: word of mouth, letters, phone calls, local radio, churches, mosques, and meetings with community members and religious leaders.

3.1.4. Training Community Service Providers

Training of community service providers cover the following:

- ◆ Community case-finding strategies.
- ◆ How to refer cases to the HP.
- ◆ How to conduct home visits for cases needing special attention.
- ◆ Acute malnutrition messages and how to disseminate them.

3.2 Implementing Community Outreach

3.2.1 Partnership with the Community

The HEW should maintain a constant and continuous dialogue with the community leaders and members and report back to the Woreda health office through the PHCU. They should involve the HDA/HDG, religious leaders, health committees, and agriculture extension workers in community outreach activities. This will increase awareness and ownership of acute malnutrition services.

How can the HEW ensure continuous dialogue with the community?

- ◆ Develop monthly plans with the HDA/HDG.
- ◆ Hold monthly meetings with the HDA/HDG to review plans and address concerns, maintain changes in behaviour, and share success stories.
- ◆ In collaboration with the HDA/HDG, organise monthly community meetings to discuss and disseminate information on acute malnutrition and other health and nutrition topics.

How can you know that there is good community involvement in acute malnutrition?

- ◆ Caregivers seek early treatment for acute malnutrition.
- ◆ A decrease in the number of patients with SAM with medical complications.
- ◆ A reduction in the default rate.

3.2.2 Case Finding and Referral

The following are strategies for case finding:

- ◆ House-to-house screening visits by HEW.
- ◆ Community screening during community health days, growth monitoring, and routine immunisation. Other available opportunities are community meetings, market days, church, general food distribution, school events, and other community-based nutrition sensitive activities.
- ◆ Screening at health facility outpatient department, inpatient wards, GMP, HIV and TB clinics, antenatal and postnatal clinics.
- ◆ Self-referrals from the communities as community members become more aware of acute malnutrition.

How should case finding be conducted?

Children 6-59 months and PLW:

- ◆ Measure MUAC
- ◆ Assess for bilateral pitting oedema

Infants under 6 months of age:

- ◆ Assess for bilateral pitting oedema
- ◆ Assess for recent weight loss or failure to gain weight.

3.2.3 Follow-up and Home Visits

The HEW should determine priority cases that need a home visit and coordinate with the HDA/HDG to monitor the patients. A home visit is critical for patients who:

- ◆ Are losing or not gaining weight, based on the action protocol (see Table 6).
- ◆ Have a deteriorating medical condition (see Table 6).
- ◆ Are not responding to treatment (see Table 6).
- ◆ Have refused referral to the SC.
- ◆ Are absent or defaulted treatment (defaulters are patients absent for 2 consecutive visits).
- ◆ Are infants < 6 months of age and have been discharged from SC.

Annex 1, the Home Visit Checklist should be used as a guide during the home visit.


3.2.4 Referral to Other Programmes and Initiatives

Families of patients with SAM and MAM, and PLW should be enrolled for PSNP. The primary caregiver of a SAM or MAM child should be exempt from public work during the treatment period. They should also be referred to other available health, WASH, food security and livelihoods interventions.

4 Management of Severe Acute Malnutrition (SAM)

This chapter describes the principles, criteria and process of managing children with SAM in the SC and OTP. The chapter also highlights on how to triage and the decision-making process for SAM case management. See Figure 2 below.

Figure 2. Decision Process for SAM Case Management



* **Medical complications include:** Poor appetite, intractable vomiting, convulsions, lethargy, not alert, unconsciousness, high fever (axillary temperature >39 °C), lower respiratory tract infection, dehydration, persistent diarrhoea, severe anaemia, hypoglycaemia, hypothermia (axillary temp. < 35 °C), severe skin lesions, eye signs of vitamin A deficiency.

4.1 Outpatient Therapeutic Programme (OTP) for the Management of SAM Without Medical Complications

OTP provides services to children with SAM with a good appetite and no medical complications. Treatment and care are provided at home with weekly follow-up visits at a nearby health facility.

Table 3. OTP Admission Criteria

- Bilateral pitting oedema + or ++
OR
 - MUAC <11.5 cm
OR
 - WFH <-3 z-score
AND
 - Appetite test passed
 - No medical complications
 - Clinically well and alert
- OR Referred from SC after stabilisation

Table 4. OTP Entry Categories

Category	Definition
New admission	New case who meets the admission criteria for OTP (see Table 3).
Readmission	Re-admitted for treatment in OTP due to relapse or a returned defaulter who meets the admission criteria.
Relapse	Cured within the past 3 months and now meets the admission criteria for OTP (see Table 3). Relapse cases require special attention, therefore should be referred to start treatment in SC.
Returned defaulters	Defaulted within the past 3 months and has returned to continue treatment in OTP. A returned defaulter should be re-admitted if they meet the admission criteria.
Transfers-in	Continuing treatment in OTP after stabilisation in SC or has moved in from another facility where s/he was receiving OTP.

4.1.2 Admission Procedure for Children 6-59 Months

STEP 1: Triage

- ◆ Identify and manage very sick patients and those referred from the community or SC first, before the health and nutrition education session is conducted.

Take Anthropometric Measurements to Confirm Nutritional Status

- ◆ Determine the age of the patient based on the health card or caregiver's recall.
- ◆ Check for presence of bilateral pitting oedema and record the findings.
- ◆ Measure MUAC and record the measurement.
- ◆ Measure and record weight.
- ◆ Measure and record height or length (if at HC level).
- ◆ Determine WFH/WFL z-score using the reference charts (if at HC level).
- ◆ If infant 0-6 months of age, assess for recent weight loss or failure to gain weight, and assess feeding according to IMNCI.
- ◆ Classify nutritional status.

Conduct an RUTF Appetite Test for Children 6-59 Months

- ◆ Conduct the RUTF appetite test in a quiet, separate area.
- ◆ Explain to the caregiver the purpose and procedure of the appetite test.
- ◆ Ask the caregiver to:
 - ◆ Wash hands with soap before feeding the RUTF.
 - ◆ Gently offer the RUTF.
 - ◆ Encourage the child to eat the RUTF without force-feeding.
 - ◆ Offer plenty of clean, safe drinking water.
- ◆ Give the caregiver one sachet of RUTF.
- ◆ Observe the child eating the RUTF.
- ◆ Determine whether he/she passes or fails the appetite test (see Table 5).

Table 5. RUTF Appetite Test Results

Pass Appetite Test	Fail Appetite Test
The child eats some of the RUTF within 30 minutes.	The child refuses to eat the RUTF after 30 minutes.

Conduct a Medical Assessment

- ◆ Conduct a medical assessment and record in the OTP Treatment and Follow-up Card. [See Annex 17.](#)
- ◆ Take the medical history, immunisation, feeding history, and assess signs and symptoms.
- ◆ Conduct a physical examination and record the findings.
- ◆ Diagnose and treat malaria according to the National Malaria Guidelines.
- ◆ Check the patient’s HIV and/or TB status if known.
- ◆ Determine whether the child needs referral to SC or treatment in the OTP. [See Table 6](#) below for the action protocol.
- ◆ Explain the findings of the medical assessment to the caregiver.

Table 6. OTP Action Protocol

Sign	Referral to SC	Follow-Up Home Visit
BILATERAL PITTING OEDEMA	Grade +++	Bilateral pitting oedema not reducing by week 3
	Severe wasting with bilateral pitting oedema	
	Develops or increases in severity	
	No reduction for 2 weeks	
	Not resolved after 3 weeks	
APPETITE / ANOREXIA	No appetite or unable to eat, fail RUTF appetite test.	Child eats < 75% of the RUTF a week by third visit
VOMITING	Intractable	General medical deterioration
TEMPERATURE	Fever: axillary temperature > 39°C	
	Hypothermia: axillary temperature < 35°C	
RESPIRATION RATE (rr)	≥ 60 respirations/minute for under 2 months	
	≥ 50 respirations/minute from 2 to 12 months	
	≥ 40 respirations/minute from 1 year to 5 years	
	≥ 30 respirations/minute for over 5 years	
	Any chest in-drawing	
ANAEMIA	Very pale (severe palmer pallor), difficulty breathing	
SUPERFICIAL INFECTION	Extensive infection requiring intravenous or intramuscular treatment	
ALERTNESS	Very weak, apathetic, unconscious	
	Fitting/convulsions	
HYDRATION STATUS	Dehydration based primarily on recent history of diarrhoea, vomiting, fever or sweating and on recent appearance of clinical signs of dehydration as reported by the caregiver or patient	
WEIGHT CHANGES		Below admission weight on week 3 (non-oedematous children)
	Weight loss for 3 consecutive weighing (non-oedematous children)	Weight loss for 2 consecutive weeks (non-oedematous children)
	Static weight for 5 consecutive weighing	Static weight for 3 consecutive weeks
GENERAL	Caregiver or patient requests SC	Returned from SC (first 2 weeks)
		Refused referral to SC
NOT RESPONDING	Patient that is not responding to treatment is referred to SC for further investigation	

Refer the Patient who Fails Appetite Test and Has Medical Complications to the SC

If the child is 6-59 months old and has medical complications or fails the appetite test, start life-saving treatment¹:

- ◆ Provide sugar water solution to avoid hypoglycaemia. Sugar water contains 10 g of sugar dissolved in 100 ml of water.
- ◆ Refer the child to the closest SC.
- ◆ Record findings and the treatment given on the Referral Slip. [See Annex 22.](#)
- ◆ Explain the following to the caregiver:
 - ◇ Severity of the condition and the need for referral to SC
 - ◇ Need to keep the child warm during transportation
 - ◇ Need to give frequent, small amounts of 10 percent sugar water solution during transportation
 - ◇ If breastfeeding, need to frequently breastfeed the child

NOTE: All infants 0-6 months of age with SAM with or without medical complications should be referred to the SC.

STEP 2: Admit Patients with Appetite and No Medical Complications to the OTP

If the child 6-59 months has no medical complications and has a good appetite, admit to the OTP:

- ◆ Explain the OTP treatment procedures and care that will be provided.
- ◆ Decide which OTP admission entry category to assign to the patient [\(see Table 4\).](#)
- ◆ Assign a registration number if the patient is a new admission.
- ◆ Give routine medicines [\(see Table 7\).](#)
- ◆ Give RUTF based on the child's weight [\(see Table 8\).](#)
- ◆ Complete the OTP Treatment and Follow-up Card.
- ◆ Register the patient in the Registration Book for SAM Treatment. [See Annex 23.](#)
- ◆ Advise the caregiver to seek medical care at the nearest health facility immediately if child becomes ill while at home or refuses to eat RUTF.

4.1.3 Medical Treatment and Care for Children 6-59 Months

Give routine medicines as indicated in [Table 7](#) below. Carefully check the records of patients who were previously treated in SC or other OTP to avoid prescribing medicines that have already been given.

¹**Medical complications include:** poor appetite, intractable vomiting, convulsions, lethargy, not alert, unconsciousness, high fever (>39 °C), lower respiratory tract infection, dehydration, persistent diarrhoea, severe anaemia, hypoglycaemia, hypothermia (<35 °C), severe skin lesions, eye signs of vitamin A deficiency.

Table 7. Routine Medication and Treatment in OTP

Name of Medication	When to Give	Age	Dose	Prescription
10% sugar water solution	During referral to SC to prevent hypoglycaemia	All ages	50 ml	Once, orally
AMOXICILLIN	On admission of all SAM patients	All ages	25 mg/kg, every 12 hours, for 5 days.	Oral: 25 mg/kg, every 12 hours, for 5 days
ANTIMALARIAL	On admission if positive malaria test	All ages	Refer to the National Malaria Guidelines	Refer to the National Malaria Guidelines
ALBENDAZOLE or MEBENDAZOLE	After 1 week for treatment in OTP	24- 59 months	Albendazole 400 mg Mebendazole 500 mg	Single dose, orally
MEASLES VACCINE	On the fourth week if the child has not yet received the measles vaccine	9–59 months	Refer to the National Expanded Programme on Immunisation (EPI) Guidelines	Refer to the National EPI Guidelines
OTHER VACCINES	Update vaccinations based on EPI schedule	All ages	Refer to the National EPI Guidelines	Refer to the National EPI Guidelines

Important Notes on Routine Medication and Treatment in OTP

HIV/AIDS and TB: If possible, offer HIV counselling and testing, and TB testing to all patients with SAM, especially if readmitted or not responding to treatment.

Antibiotic treatment with Amoxicillin: If the patient is HIV+ or HIV-exposed and taking Cotrimoxazole prophylaxis, Amoxicillin should also be given. Medical treatment and care should follow the National Guidelines for Comprehensive HIV Prevention, Treatment and Care.

Vitamin A: Do not give an additional high dose of vitamin A to children with SAM in the OTP as RUTF contains an adequate amount of vitamin A.

- ◆ A high dose (50,000 IU, 100,000 IU or 200,000 IU, depending on age) of vitamin A should be given on admission only if the therapeutic foods given are not fortified as recommended in WHO specifications and if vitamin A is not part of other daily supplements and was not given to the child in the past one month. (Note: RUTF is fortified).
- ◆ A high dose of vitamin A should be given on day 1 to all patients with SAM and eye signs of vitamin A deficiency (e.g., night blindness, Bitot’s spots, corneal ulceration) and to all patients with SAM and recent measles (now or in the past 3 months). The second and third doses should be given on day 2 and day 15, irrespective of the type of therapeutic food they are receiving.
- ◆ All patients who show signs of vitamin A deficiency should be referred for treatment in SC.

Iron and Folic Acid: Iron and folic acid should not be given routinely to children in OTP as RUTF contains iron and folic acid. If anaemia is identified, it should be treated according to IMNCI/ICCM-CBNC guidelines, and treatment should begin after 14 days of SAM treatment. If severe anaemia, refer for treatment in SC. **Malaria testing and treatment should be done before the iron and folic acid treatment is given.**

Zinc: Do not give zinc supplements to treat diarrhoea in SAM patients in the OTP. RUTF contains adequate amounts of zinc.

ORS: ORS contains high sodium and is inappropriate for patients with SAM. ORS is only used if the SAM patient has profuse watery diarrhoea, or AWD. Patients who require treatment for dehydration should be referred for SC. In SC, patients with dehydration are treated with ReSoMal. The use of ReSoMal is inappropriate in an OTP setting.

4.1.4 Nutrition Rehabilitation for Children 6-59 Months

- ◆ Use the reference **Table 8** below to determine the amount of RUTF to give at each weekly visit, based on the patient's weight.

Table 8. RUTF Reference Table (92 g Sachet Containing 500 Kcal)

Weight of Child (kg)	Sachets per Day	Sachets per Week
3.5 - 3.9	1½	11
4.0 – 5.4	2	14
5.5 – 6.9	2½	18
7.0 – 8.4	3	21
8.5 – 9.4	3½	25
9.5 – 10.4	4	28
10.5 – 11.9	4½	32
≥ 12	5	35

-
- ◆ Explain the following on the feeding procedure:
 - ◆ **Do not give RUTF to infants 0-6 months.** They should be exclusively breastfed.
 - ◆ If the child is 6-23 months, continue to breastfeed on demand. Mother's milk is best for infants and young children.
 - ◆ Feed small amounts of RUTF until the allocated daily ration is finished, and before eating any other food (apart from breast milk).
 - ◆ Encourage the child to eat as often as possible (every 3 hours during the day).
 - ◆ If the child is breastfeeding, offer breast milk on demand and before feeding RUTF.
 - ◆ Offer plenty of safe drinking water while eating RUTF.
 - ◆ Do not mix RUTF with liquids as this might cause bacterial growth.

4.1.5 Monitoring During OTP Follow-Up Visits

- ◆ Monitor the following parameters and record information on the OTP Treatment and Follow-up Card at each follow-up visits.
 - ◆ MUAC
 - ◆ Weight
 - ◆ Height/length (if at HC level, measure once a month)
 - ◆ WFH/WFL z-score (if at HC level, check on admission, every month, and on discharge)
 - ◆ Degree of bilateral pitting oedema (0, +, ++, +++)
 - ◆ Weight loss (patients who lose weight or have weight fluctuations should receive a home visit and/or further medical examination)
 - ◆ Clinical signs: body temperature, respiration rate, dehydration, anaemia, and skin infection
 - ◆ Standard medical history: stool, vomiting, fever, and cough
 - ◆ Appetite test
 - ◆ Any illness suffered since the last visit
 - ◆ Any action taken, or medication given in response to a health condition
- ◆ Repeat the RUTF feeding procedure on every follow-up visit and ask the caregiver to repeat back.
- ◆ Inform the caregiver on the treatment progress.
- ◆ Conduct individual and/or group counselling on health, WASH, and IYCF.

-
- ◆ Indicate any necessary follow-up actions on the OTP Treatment and Follow-up Card, such as:
 - ◆ Home visits for patients requiring special attention. See Table 6 for the action protocol.
 - ◆ Referral to the SC.

Prioritize Patients for a Home Visit

Conduct a home visit if the patient:

- ◆ Is losing or not gaining weight
- ◆ Has a deteriorating medical condition
- ◆ Is not responding to treatment
- ◆ Has refused SC
- ◆ Is absent or defaulted treatment (defaulters are patients absent for 2 consecutive visits)

Referral to SC Based on the Action Protocol

- ◆ Refer patients with a poor appetite, deteriorating nutritional status, and/or a medical complication. See Table 6 for the action protocol.
- ◆ Issue a Referral Slip to the patient referred for treatment in the SC.

Failure to Respond to Treatment

It is recommended that patients who are referred to SC due to failure to respond to treatment be tested for HIV and TB which are common underlying causes of acute malnutrition.

The following are other frequent causes of failure to respond to treatment:

- ◆ Inappropriate evaluation of the patient's health condition or missed medical complication
- ◆ Non-adherence to the RUTF protocol
- ◆ Non-adherence to the routine medication protocol
- ◆ Inadequate guidance provided for home care
- ◆ Inappropriate frequency of visits to the health facility and reception of RUTF
- ◆ Inadequate intake or sharing of RUTF and/or medicines with other family members

4.1.6 Discharge Criteria and Exit Categories for Children 6-59 Months

Table 9. OTP Discharge Criteria

Note: The same anthropometric indicator that is used to identify and confirm SAM on admission should be used to determine recovery and discharge from treatment

If admitted with bilateral pitting oedema, discharge cured when:

- No bilateral pitting oedema for 2 consecutive visits
- **AND** MUAC \geq 12.5 cm or WFH/WFL \geq -2 z-score
- **AND** Clinically well and alert

If admitted based on MUAC, discharge cured when:

- MUAC \geq 12.5 cm
- **AND** No bilateral pitting oedema
- **AND** Clinically well and alert

If admitted based on WFH/WFL, discharge cured when:

- WFH/WFL \geq -2 z-score
- **AND** No bilateral pitting oedema
- **AND** Clinically well and alert.

Table 10. OTP Exit Categories

Category	Definition
Cured	Has reached the discharge criteria for SAM treatment (see Table 9).
Died	Dies while receiving treatment in the OTP.
Defaulted	Absent for two consecutive visits. Default should be confirmed.
Non-responder	Does not reach the SAM discharge criteria after 16 weeks (4 months) in treatment.
Transferred-out	Condition has deteriorated or not responding to treatment according to action protocol and referred for treatment in the SC, or moved out to receive OTP in another facility.

4.1.7 Discharge Procedure for Children 6-59 Months

- ◆ Give feedback to the caregiver on the final treatment outcome.
- ◆ Ensure that the caregiver understands how to use continuing medications.
- ◆ Give the final on week supply of RUTF ration.
- ◆ Record the exit category (see Table 10) on the Registration Book for SAM and the OTP Treatment and Follow-up Card.
- ◆ Counsel on WASH, good nutrition, and IYCF practices.
- ◆ Advise the caregiver to immediately go to the nearest health facility if they experience any signs of deterioration.
- ◆ Refer the caregiver to PSNP and other complementary health and nutrition services.

4.2 Stabilisation Centre (SC) for the Management of SAM With Medical Complications

4.2.1 Admission Criteria and Entry Categories for Children 6-59 Months

Table 11. SC Admission Criteria

- ◆ Bilateral pitting oedema +++
OR
 - ◆ Any grade of bilateral pitting oedema combined with severe wasting (MUAC <11.5 cm or WFH < -3 z score)
OR
 - ◆ Bilateral pitting oedema +, ++
OR
 - ◆ Severe wasting (MUAC <11.5 cm or WFH < -3 z score)
WITH any of the following medical complications:
 - ◆ Poor appetite
 - ◆ Intractable vomiting
 - ◆ Convulsions
 - ◆ Lethargy, not alert
 - ◆ Unconsciousness
 - ◆ High fever (axillary temperature > 39°C)
 - ◆ Lower respiratory tract infection
 - ◆ Dehydration
 - ◆ Persistent diarrhoea
 - ◆ Severe anaemia
 - ◆ Hypoglycaemia
 - ◆ Hypothermia (axillary temperature < 35°C)
 - ◆ Severe skin lesions
 - ◆ Eye signs of vitamin A deficiency
- OR Referred from outpatient care according to the action protocol

Table 12. SC Entry Categories

Category	Definition
New admissions	New case who meets the admission criteria for SC (see Table 11).
Readmission	Re-admitted for treatment in SC due to relapse or a returned defaulter who meets the admission criteria.
Relapse	Cured within the past 3 months and now meets the admission criteria for SC (see Table 11).
Returned defaulters	Defaulted within the past 3 weeks and has returned to continue treatment in SC. A returned defaulter should be re-admitted if they meet the admission criteria.
Transfers-in	Referred to the SC because condition deteriorated in OTP or has moved in from another facility where s/he was receiving care in SC.

4.2.2 Admission Procedure

STEP 1: Triage

Follow Emergency Triage Assessment and Treatment (ETAT) principles. Patients with SAM with medical complications should be identified as having priority signs that require prompt assessment and treatment. Assess and treat the following:

- ◆ Obstructed breathing
- ◆ Severe respiratory distress
- ◆ Central cyanosis
- ◆ Shock (cold hands, capillary refill time longer than 3 seconds, fast heart rate with weak pulse, and low or unmeasurable blood pressure)
- ◆ Unconsciousness
- ◆ Convulsions
- ◆ Dehydration
- ◆ Severe anaemia
- ◆ Hypoglycaemia
- ◆ Hypothermia
- ◆ Congestive heart failure

If any of the above signs are present, begin immediate emergency treatment.

STEP 2: Take Anthropometric Measurements to Confirm Nutritional Status

- ◆ Determine the age of the patient based on the health card or caregiver's recall.
- ◆ Check for presence of bilateral pitting oedema and record the findings.
- ◆ Measure MUAC and record the measurement.
- ◆ Measure and record weight.
- ◆ Measure and record height or length.
- ◆ Determine WFH/WFL z-score using the reference charts.
- ◆ Classify nutritional status.

STEP 3: Conduct Medical Assessment

- ◆ Take a medical history to determine risk factors for SAM and record the following information on the SC Multi-chart. [See Annex 18.](#)
- ◆ Take a feeding history and record findings in the SC Multi-chart.
- ◆ Conduct a general physical examination and examine for complications of SAM.
- ◆ Diagnose and treat malaria according to the National Malaria Guidelines.
- ◆ Offer HIV counselling and testing services; if positive, link the patient to HIV treatment and care. Refer to the current National Guidelines for Comprehensive HIV Prevention, Treatment and Care.
- ◆ Check:
 - ◆ Blood glucose
 - ◆ Haemoglobin
 - ◆ Urine analysis (to rule out nephrotic syndrome in oedematous patients)

STEP 4: Admit and Initiate Medical and Nutrition Treatment in SC

- ◆ Explain to the caregiver the SC treatment procedures and care that will be provided.
 - ◆ Initiate feeding and medical treatment immediately.
 - ◆ Complete records as follows:
 - ◆ Assign a registration number if the patient is a new admission.
 - ◆ Decide the SC entry category to assign the patient. [See Table 12.](#)
 - ◆ Complete the SC Multi-chart.
 - ◆ Register the patient in the Registration Book for SAM Treatment.
-

4.2.3 Treatment and Care of Children 6-59 Months

Treatment and care of patients with SAM with medical complications is divided into the following phases:

- ◆ Stabilisation phase in which life-threatening problems are identified and treated, specific deficiencies are corrected, metabolic abnormalities are reversed, and feeding is established using F-75.
- ◆ Transition which prepares the patient for the rehabilitation phase. RUTF (or F-100) is gradually introduced and the patient's condition carefully monitored.
- ◆ Rehabilitation phase when intensive feeding is given to recover most of the lost weight, emotional and physical stimulation are increased. In most cases, rehabilitation will take place in OTP using RUTF. There are a few exceptions where the patient cannot be safely transferred to OTP:
 - ◆ Patients who do not tolerate RUTF, in this case F-100 is used for rehabilitation.
 - ◆ Patients who have no access to an OTP near their homes.
 - ◆ Patients whose caregiver refuses OTP, despite being counselled.

4.2.3.1 Treat and Prevent Hypoglycaemia

Hypoglycaemia is a low level of glucose in the blood. In a child with SAM, the level considered low is than < 3 mmol/litre (or < 54 mg/dl). Hypoglycaemia and hypothermia (low temperature) usually occur together and are signs of infection. Check for hypoglycaemia whenever hypothermia (axillary temperature $< 35.0^{\circ}$ C or rectal temperature $< 35.5^{\circ}$ C) is found. Frequent feeding is important in preventing both conditions.

Prevention

- ◆ Feed F-75 every 2 hours, starting straight away, if possible.
- ◆ Always give feeds throughout the day and night.

Treatment

If the patient is conscious and a glucose test shows < 3 mmol/L or 54 mg/dl, give:

- ◆ A 50 ml bolus of 10% glucose or 10% sucrose or 10% dextrose solution (i.e., 37.5 ml of distilled water with 12.5 ml of 40% dextrose) orally or by NGT, then feed F-75 every 30 minutes for 2 hours, (giving 1/4 of the 2-hour F-75 feed each time).
- ◆ Give antibiotics (see Table 13).
- ◆ Feed F-75 every 2 hours, day and night.

If the patient is unconscious, lethargic or convulsing, give:

- ◆ Intravenous (IV) sterile 10 % glucose solution (5 ml/kg of body weight), followed by 50 ml of 10% glucose or sucrose or dextrose solution by NGT, then feed F-75 every 30 minutes for 2 hours, giving 1/4 of the 2-hour feed each time.
- ◆ Give antibiotics (see Table 13).
- ◆ Feed F-75 every 2 hours, day and night.

Monitor

- ◆ Blood glucose:
 - ◇ If low, repeat the glucose test after two hours, taking blood from the child's finger or heel. Once treated, most patients stabilise within 30 minutes.
 - ◇ If blood glucose falls to < 3 mmol/L, give a further 50 ml of 10% glucose or sucrose solution and continue feeding every 30 minutes until stable.
 - ◇ If blood glucose is 3 mmol/L or higher, change to 2-hourly feeds of F-75. If still low, make sure antibiotics and F-75 have been given. Keep giving F-75 every half-hour.
- ◆ Axillary temperature: If < 35.0° C, repeat the glucose test.
- ◆ Level of consciousness: If this deteriorates, repeat the glucose test.

Note: If you are unable to test for blood glucose levels, assume that all children with SAM have hypoglycaemia and treat accordingly.

4.2.3.2 Treat and Prevent Hypothermia

Hypothermia is low body temperature. A SAM patient is hypothermic if the axillary temperature is below 35°C or if the rectal temperature is below 35.5°C.

Prevention

- ◆ Feed F-75 every 2 hours, starting straight away.
- ◆ Always give feeds throughout the day and night.
- ◆ Keep the patient covered and away from drafts.
- ◆ Warm your hands before touching the patient (both health service providers and caregivers).
- ◆ Keep the child dry and promptly change wet nappies, clothes, and bedding.
- ◆ Avoid exposure (e.g., bathing, prolonged medical examinations).
- ◆ Let the child sleep with the mother/caregiver at night for warmth.
- ◆ Maintain room temperature between 28° C and 32° C.

Treatment

If the axillary temperature is < 35.0° C or rectal temperature is < 35.5° C:

- ◆ Feed F-75 straight away (or start rehydration, if needed).
- ◆ Re-warm the patient: Either clothe the patient (including head), cover with a warmed blanket, and place a heater or lamp nearby (do not use a hot water bottle), or put the child on the mother's bare chest (skin to skin) and cover both of them (kangaroo technique).
- ◆ Give antibiotics (see Table 13).

Monitor

- ◆ Body temperature every 30 minutes for the first 2 hours
- ◆ Body temperature: During re-warming, take temperature every 2 hours until it rises to > 36.5° C (take every 30 minutes if heater is used).
- ◆ Ensure the patient is always covered, especially at night.
- ◆ Feel for warmth.
- ◆ Blood glucose level: Check for hypoglycaemia whenever hypothermia is diagnosed.

4.2.3.3 Treat and Prevent Dehydration

- ◆ Dehydration is difficult to diagnose in patients with SAM. Clinical signs associated with dehydration in those well-nourished (dry mouth, sunken eyes, skin lying in folds, etc.) are often present in SAM patients without dehydration.
- ◆ Ask if the patient had watery diarrhoea or vomiting, accompanied by recent changes in appearance; if yes, assume dehydration and give ReSoMal. ReSoMal is a modification of ORS, it contains less sodium, more sugar, and more potassium than ORS and is intended for SAM children with diarrhea.
- ◆ If the patient is diagnosed with profuse watery diarrhea or AWD, ORS is recommended instead of ReSoMal. See Annex 2 on the management of AWD and SAM.

Prevention

To prevent dehydration when a patient has continuing watery diarrhoea:


- ◆ Keep feeding with F-75.
 - ◆ Replace the approximate volume of stool losses with ReSoMal.
 - ◆ Give ReSoMal after each watery stool: for children without bilateral pitting oedema, 50–100 ml if child is under 2 years, and 100-200ml if child is above 2 years.
 - ◆ If the child has bilateral pitting oedema, give 30 ml of ReSoMal per each watery stool.
 - ◆ NOTE: It is common for SAM children to pass many small, unformed stools. These should not be confused with profuse watery stools and do not require fluid replacement.
 - ◆ If the child is breastfed, encourage the caregiver to continue breastfeeding.
-

Treatment

- ◆ Give ReSoMal 5 mls/kg every 30 minutes for the first 2 hours.
- ◆ Then, if still dehydrated, give ReSoMal 5–10 ml/kg per hour. Alternate with F-75, up to a maximum of 10 hours.
- ◆ Encourage the mother/caregiver to give the fluid slowly and to persist, even if the child is slow to take the fluids. Give the mother/caregiver only the amount of fluid required for the next hour.
- ◆ If the patient is refusing or vomiting, insert NGT and commence fluids via NGT.
- ◆ Length of rehydration is dependent on the patient’s thirst and fluid loss from vomiting and stools.
- ◆ Do not treat dehydration with intravenous fluids because children can become overloaded with fluid very quickly, which is very dangerous and can lead to heart failure and death.
- ◆ If breastfeeding, encourage the caregiver to continue.

During treatment, rapid respiration and pulse rates should slow down and the child should begin to pass urine.

Figure 3. Treatment of Dehydration



Monitor Rehydration Progress


Observe the patient every 30 minutes for 2 hours, then hourly for the next 6–12 hours, recording:

- ◆ Pulse rate
- ◆ Respiratory rate
- ◆ Urine frequency
- ◆ Stool/vomit frequency
- ◆ Signs of improvement:
 - ◇ Decreasing pulse and respiration rate
 - ◇ Reduced thirst
 - ◇ Increasing urinary output
 - ◇ Patient is more alert
- ◆ Signs of over-hydration
 - ◇ Increased heart rate (by 25 beats a minute)
 - ◇ Increased respiration (by at least five breaths a minute)
 - ◇ Signs of increased oedema (such as puffy eyelids)
 - ◇ Prominent neck veins and enlargement of the liver

Continued rapid breathing and pulse during rehydration suggest coexisting infection or over-hydration. If there are signs of over-hydration, stop ReSoMal immediately and reassess in one hour.

NOTE: Low blood volume can coexist with oedema. Do not use an IV for rehydration except in cases of shock, and then do so with care, infusing slowly to avoid flooding the circulation and overloading the heart.

Figure 4. Monitoring Patients on ReSoMal



4.2.3.4 Correct Electrolyte Imbalance

- ◆ Children with SAM have excess body sodium, even though plasma sodium may be low; giving high sodium loads will cause death. They are also deficient in potassium and magnesium and these deficiencies may take at least 2 weeks to correct. Oedema is partly due to these imbalances. Do not treat oedema with a diuretic.
- ◆ Therapeutic foods (F-75, F-100, and RUTF) and ReSoMal contain extra potassium and magnesium.

4.2.3.5 Treat and Prevent Infection

In a patient with SAM, typical signs of infection, such as fever, are often absent and infections are often hidden. Therefore, give a broad-spectrum antibiotic routinely on admission. [Table 13](#) below summarises the choice of broad-spectrum antibiotics for patients with SAM. Also [see Annex 3](#) for the antibiotics and formulations.

Table 13. Choice of Broad-Spectrum Antibiotics in SC

IF:	GIVE:		
NO MEDICAL COMPLICATIONS	Amoxicillin oral: 25 mg/kg every 12 hours for 5 days or until referral to OTP		
MEDICAL COMPLICATIONS (shock, hypoglycaemia, hypothermia, dermatosis with raw skin/fissures, respiratory or urinary tract infections, or lethargic/sickly appearance)	<p>*Gentamicin IV or IM (5 mg/kg) once daily for 7 days, plus:</p> <table border="1"> <tr> <td>Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days</td> <td>Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days</td> </tr> </table>	Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days	Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days
Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days	Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days		
Resistance to Amoxicillin and Ampicillin and Presence of Medical Complications	<p>In the case of sepsis or septic shock, give: IV/IM ceftriaxone (children or infants over 1 month of age (50 mg/kg every 8 to 12 hours) + oral ciprofloxacin (5 to 15 mg/kg 2 times per day)</p> <p>If suspected staphylococcal infection, add: IV/IM cloxacillin (12.5–50 mg/kg/dose four times a day, depending on the severity of the infection).</p>		
Specific Infection Requires an Additional Antibiotic	Add specific antibiotic as per the Ethiopia Standard Treatment Guidelines.		
If HIV-positive or Child who is HIV Exposed.	Cotrimoxazole oral according to the National Guidelines for Comprehensive HIV Prevention, Treatment and Care.		

*If the child is not passing urine, gentamicin may accumulate in the body and cause deafness. Do not give the second dose until the child is passing urine.

**If amoxicillin is not available, give ampicillin, 50 mg/kg orally every 6 hours for 5 days.

4.2.3.6 Treatment of SAM Complications and Associated Conditions

Shock

Shock from dehydration and sepsis are likely to coexist in children with SAM. They are difficult to differentiate on clinical signs alone. Children with dehydration will respond to IV fluids. Those with septic shock and no dehydration will not respond. The amount of fluid given is determined by the child's response. Over hydration must be avoided.

To start treatment:

- ◆ Give oxygen.
- ◆ Give sterile 10% glucose (5 ml/kg) by IV.
- ◆ Give IV fluid at 15 ml/kg over 1 hour. Use Ringer's lactate with 5% dextrose; or 0.45% half-normal saline with 5% dextrose.
- ◆ Measure and record pulse and respiration rates every 10 minutes.
- ◆ Give antibiotics (see Table 13).

If there are signs of improvement (pulse and respiration rates are lower after 1 hour):

- ◆ Repeat IV 15 ml/kg over 1 hour, then switch to oral or NGT rehydration with ReSoMal, 10 ml/kg/h for up to 10 hours. (Leave IV in place in case required again).
- ◆ Give ReSoMal in alternate hours with F-75, then continue feeding with starter F-75.

If the child fails to improve after the first hour of treatment (15 ml/kg), assume that the child has septic shock.

In this case:

- ◆ Give maintenance IV fluids (4 ml/kg/h) while waiting for blood,
- ◆ When blood is available, transfuse fresh whole blood at 10 ml/kg slowly over 3 hours.
- ◆ Then, begin feeding with F-75.

If the child gets worse during treatment (breathing increases by 5 breaths or more/min and pulse increases by 25 or more beats/min): stop the infusion to prevent the child's condition worsening.

Severe Anaemia

Anaemia is a serious problem if it is very severe. Give a blood transfusion if the haemoglobin (Hb) concentration is < 4 g/dl or if Hb concentration is 4-6 g/dl and there are signs of congestive heart failure.

- ◆ Stop oral intake and V during the blood transfusion.
- ◆ Give Furosemide 1 mg/kg intravenously at the start of the blood transfusion.
- ◆ Give a blood transfusion of whole fresh blood (10 ml/kg) slowly over 3 hours (or 5-7 ml/kg packed red cells if there are signs of congestive heart failure).

- ◆ Wait 3 hours after the transfusion before feeding.
- ◆ Do not give a second transfusion, even if the haemoglobin is still low, for at least 3 days.
- ◆ Do not give iron in the stabilisation phase or during transition.

Measles

Give measles vaccine according to the Expanded Programme on Immunisation (EPI) schedule. If the child has an active case of measles or has had measles recently (in the past 3 months), give a high dose of vitamin A on days 1, 2, and 15, following the dosages in [Table 14](#).

Table 14. Vitamin A Treatment Dosages

Age	Oral Dose
< 6 months	50,000 IU (15,000 µg)
6–11 months	100,000 IU (30,000 µg)
≥ 12 months	200,000 IU (60,000 µg)

Eye Signs of Vitamin A Deficiency

- ◆ Corneal ulceration due to vitamin A deficiency requires emergency treatment and care.
- ◆ Put 1 drop of Atropine into the affected eye three times daily to relax the eye and prevent the lens from being pushed out.
- ◆ Give a high dose of vitamin A on days 1, 2, and 15, following dosages in [Table 14](#).
- ◆ Cover the affected eye with a damp gauze pad (dampen with 0.9% saline) and bandage to hold the pad in place.
- ◆ If necessary, put mittens or bandages on the child’s hands to prevent him/her from touching his/her eyes.
- ◆ If there is pus or inflammation, use Chloramphenicol or Tetracycline eye drops 4 times daily for 7–10 days.

NOTE: Children with vitamin A deficiency are likely to be photophobic and will keep their eyes closed. It is important to examine the eyes very gently to prevent corneal rupture.

Dermatosis (Skin Lesions)

Signs:

- ◆ Hypo-pigmentation or hyper-pigmentation
- ◆ Desquamation
- ◆ Ulceration (spreading over limbs, thighs, genitalia, groin, and/or behind the ears)
- ◆ Exudative lesions (resembling severe burns), often with secondary infections, including candida

Treatment:

- ◆ Protect skin from further damage by handling the patient gently. Infections enter the body easily through broken skin or fissures.
- ◆ If the child has mild or moderate dermatosis, use regular soap for bathing.
- ◆ If severe dermatosis, bathe for 10-15 minutes/day in 1% potassium permanganate solution. This dries the lesions, helps prevent loss of serum, and inhibits infection. If the child has severe dermatosis but is too sick to be bathed, dab 1% potassium permanganate solution on the lesions and dress oozing areas with gauze to keep them clean.
- ◆ If 1% potassium permanganate is not available, dress with silver sulfadiazine impregnated tulle dressings or cream (1%) once per day; if unavailable, apply barrier cream (10% zinc oxide ointment, castor oil ointment, petroleum jelly, or gentian violet paint). Use a different tube of ointment for each child to avoid spreading infection.
- ◆ At night and in cold conditions, dress raw areas with sterile paraffin gauze.
- ◆ For sore areas around the perineum, leave nappies off and keep the area dry. Apply nystatin cream twice daily if the diaper area is colonized with candida.

Parasitic Worms

Give Albendazole at the start of rehabilitation (on day 7), when the patient is being discharged to OTP. If Albendazole is not available, give Mebendazole (see Table 15).

Table 15. Albendazole or Mebendazole

AGE	ALBENDAZOLE	MEBENDAZOLE
< 2 year	None	None
≥ 2 years	400 mg single dose	500 mg single dose

Diarrhoea

- ◆ Diarrhoea is common in SAM children, but it should subside during the first week of treatment with cautious feeding. In the rehabilitation phase, loose, poorly formed stools are no cause for concern, if weight gain is satisfactory.
- ◆ Mucosal damage and giardiasis are common causes of continuing diarrhoea. Where possible examine the stools by microscopy. If cysts or trophozoites of giardia lamblia are found, give:
 - ◆ Metronidazole (7.5 mg/kg 8-hourly for 7 days)
 - ◆ Treat with metronidazole if stool microscopy cannot be undertaken or if there is clinical suspicion of giardiasis.

Oral Thrush

Treat oral thrush with oral nystatin suspension (100,000 IU/ml). Give 1–2 ml 4 times a day for 7 days.

Malaria

Treat Malaria according to the National Malaria Guidelines.

HIV/AIDS and TB

See chapter 6 on the management of acute malnutrition in the context of HIV/AIDS and TB.

4.2.3.7 Correct Micronutrient Deficiencies

- ◆ Patients with SAM have vitamin and mineral deficiencies (especially potassium and magnesium). Although anaemia is common, do not give iron in the stabilisation phase or during transition because giving iron can make infections worse.
- ◆ Vitamins and minerals, including vitamin A, folic acid, zinc, and copper, are present in therapeutic food (F-75, F-100 and RUTF) that complies with WHO specifications, therefore it is not necessary to give additional doses of these micronutrients.
- ◆ If the patient has eye signs of vitamin A deficiency (e.g., corneal clouding, corneal ulceration, Bitot's spots), or if the patient has recent measles (now or in the past 3 months), give oral vitamin A on days 1, 2, and 15. See Table 14 for the dosage to give.

4.2.4 Feeding Children 6-59 Months

Stabilisation Phase

In the stabilisation phase, a cautious approach is required because of the patient's fragile physiological state and reduced homeostatic capacity. Feeding should be started as soon as possible after admission and should be designed to provide just enough energy and protein to maintain basic physiological processes.

- ◆ Give small, frequent feeds of F-75 every 2 hours, orally or by NGT.
- ◆ F-75 contains 75 kcal and 0.9 g of protein per 100 ml
- ◆ If breastfed child, encourage the child to continue breastfeeding.

Preparation of F-75 Feeds using the New Tin Formulation

1. Boil water to make it safe for drinking.
2. Ensure that the water temperature is not below 70°C (i.e., cooled for not less than 3-5 minutes after boiling).
3. Add the water to the F-75 therapeutic milk powder. [See Table 16.](#)
4. Whisk the mixture vigorously until the powder dissolves in the water.
5. Cool the prepared milk to feeding temperature before administering

Table 16. Preparation of F-75 Feeds using the New Tin Formulation

White Scoops of F-75	Water (ml)	*Volume of F-75 Milk (ml)
1	25	28
2	50	56
3	75	84
4	100	112
5	125	140
6	150	168
7	175	196
8	200	224
9	225	252
10	250	280
20	500	560
Tins of F-75	Water (ml)	*Volume of F-75 Milk (ml)
1 Tin (400 grams)	2200	2480
2 Tins (800 grams)	4400	4960

***Note** that the F-75 milk volumes provided are estimates; the health care provider should measure the amount of milk to feed the child based on his/her body weight.

Quantity of F-75 to Feed

The amount of F-75 per feed is calculated based on the patient's admission weight. [Annexes 4 and 5](#) provide the recommended amount of F-75 for patients with severe wasting and those with severe bilateral pitting oedema (+++). Use the day 1 weight to determine how much to give, even if the patient loses or gains weight in this phase.

Feeding Procedure in Stabilisation Phase

Feed by cup and saucer. Only feed with an NGT when the patient is unable to take enough F-75 by mouth. Enough is defined as intake of 80% of the milk. An NGT should be used if the child:

- ◆ Takes less than 80% of the prescribed diet on two consecutive feeds during stabilisation
- ◆ Has pneumonia (rapid respiration rate) and difficulty swallowing
- ◆ Has painful lesions/ulcers of the mouth
- ◆ Has a cleft palate or other physical deformity
- ◆ Is very weak and shows difficulty remaining conscious

The NGT should only be used in the stabilisation phase. [See Annex 6](#) on how to insert an NGT.

Monitoring during Stabilisation Phase

- ◆ Measure and record weight on the SC Multi-chart and plot it on the chart each day.
- ◆ Assess and record the degree of oedema (0, +, ++, +++) each day.
- ◆ Measure MUAC on admission, and then once per week.
- ◆ Monitor body temperature, pulse, and respiration every 4 hours.
- ◆ Assess and record any standard clinical signs (stools, vomiting, dehydration, cough, skin conditions, and perianal lesions) daily.
- ◆ Note and record in the patient record each day whether the patient is absent, vomits, or refuses a feed, and whether the patient is fed by NGT or is given an IV infusion or blood transfusion.

NOTE: During stabilisation, diarrhoea should gradually diminish, and children with bilateral pitting oedema should start losing weight. If diarrhoea continues despite cautious re-feeding or worsens substantially, re-evaluate the patient.

Transition

Use the following criteria to assess a patient's readiness to transition from the stabilisation phase:

- ◆ Appetite has returned (i.e., easily finishes all F-75 milk).
- ◆ Subsiding bilateral pitting oedema (e.g., severe oedema [+++]) has been reduced to at least moderate oedema [++]).
- ◆ No serious medical problems, such as vomiting, watery diarrhoea, dehydration, nasogastric feeding, respiratory distress, or any complication that requires IV infusion, are present.
- ◆ Transition Using RUTF
- ◆ During this period, introduce RUTF gradually alongside F-75. Some children may initially refuse the RUTF; continue to offer RUTF at every feed until they begin to eat the prescribed amounts.

The recommended energy intake during transition is 100-135 kcal/kg bodyweight/day.

RUTF Quantities

- ◆ The full day's amount of RUTF should be given to the caregiver and the amount taken should be checked before each feeding time.
- ◆ When the patient is taking more than 75% of the daily prescribed amount of RUTF, he/she should be referred to OTP and continue treatment at home. [See Table 17](#) for RUTF to give per day, according to the patient's body weight and the 75% daily ration.

Table 17. RUTF Reference Table (92 g Sachet Containing 500 Kcal), with the Amounts for Transition in the SC

Weight (kg)	Sachets per Day	75% of Daily Ration	Sachets per Week
3.5 - 3.9	1 ^{1/2}	1 ^{1/4}	11
4.0 - 5.4	2	1 ^{1/2}	14
5.5 - 6.9	2 ^{1/2}	2	18
7.0 - 8.4	3	2 ^{1/2}	21
8.5 - 9.4	3 ^{1/2}	2 ^{3/4}	25
9.5 - 10.4	4	3	28
10.5 - 11.9	4 ^{1/2}	3 ^{1/2}	32
≥ 12	5	4	35

Feeding Procedure during Transition

- ◆ Provide RUTF to the caregiver.
- ◆ Do not give RUTF to infants 0-6 months.
- ◆ Encourage the patient to have small, frequent RUTF feeds every 4 hours (six times per day).
- ◆ Breastfed children should be offered breast milk on demand before being fed RUTF.
- ◆ Offer clean, safe drinking water during and after the RUTF feed.
- ◆ Every 3–4 hours, provide the same amount of F-75 therapeutic milk as in the stabilisation phase.
- ◆ When the patient finishes 50 percent of the RUTF, reduce the volume of F-75 provided by 50 percent.
- ◆ Stop providing F-75 when the patient can finish 75–100 percent of the daily RUTF ration.

Transition Special Cases Who Cannot Tolerate RUTF

In rare cases, some patients are not able to consume RUTF, such as those with problems with swallowing due to severe mouth sores, cleft palate, or neurological problems such as cerebral palsy. In these circumstances, the patient should transition from F-75 to F-100. Ongoing counselling should also be given to caregivers to explain the patient's condition and the implications for feeding.

Quantities of F-100 During Transition

- ◆ The volume of feeds remains the same as in the stabilisation phase.
- ◆ Give 130 ml of F-100 (150 kcal) per kg body weight per day.
- ◆ Use the F-100 reference table (see Annex 7) for the volume of F-100 to give per feeding, according to the patient's body weight.

Preparation of F-100 using the New Tin Formulation

1. Boil water to make it safe for drinking.
2. Ensure that the water temperature is not below 70°C (i.e., cooled for not less than 3-5 minutes after boiling).
3. Add the water to the F-100 therapeutic milk powder. See Table 18.
4. Whisk the mixture vigorously until the powder dissolves in the water.
5. Cool the prepared milk to feeding temperature before administering.

Table 18. Preparation of F-100 Feeds using the New Tin Formulation

Blue Scoops of F-100	Water (ml)	*Volume of F-100 Milk (ml)
1	25	29
2	50	58
3	75	87
4	100	117
5	125	145
6	150	174
7	175	203
8	200	234
9	225	261
10	250	290
20	500	580
Tins of F-100	Water (ml)	*Volume of F-100 Milk (ml)
1 Tin (400 grams)	2200	2158
2 Tins (800 grams)	4400	4316

***Note** that the F-100 milk volumes provided are estimates; the health care provider should measure the amount of milk to feed the child based on his/her body weight.

Give the Feed based on the Child's Body Weight

The procedures and timing of F-100 feeds in the transition phase are the same as the F-75 feeds in the stabilisation phase (see "Feeding Procedure in the Stabilisation Phase" on page 62). Breastfed children should be offered breast milk on demand before being fed F-100. Patients should never be force-fed.

Monitoring during Transition

Monitor and record the following parameters on the SC Multi-chart:

- ◆ Weight
- ◆ Degree of oedema (0, +, ++, +++)
- ◆ Body temperature, pulse, and respiration
- ◆ Standard clinical signs, such as stool, vomiting, dehydration, cough, and respiration.
- ◆ MUAC (measure and record each week)
- ◆ Other relevant data (e.g., absences, refusal of feed)
- ◆ Mood or smile

Progression from the Transition Phase

- ◆ Recovering patients can progress to OTP. Only those who cannot eat RUTF remain in SC. Patients experiencing certain complications should be returned to the stabilisation phase.

Criteria to Move Back from the Transition Phase to the Stabilisation Phase

- ◆ The patient should be moved back to the stabilisation phase if there is:
- ◆ Weight gain of more than 10 g/kg/day in association with an increase in respiratory rate (indicative of excess fluid retention)
- ◆ Increasing or developing oedema
- ◆ A rapid increase in the size of the liver
- ◆ Any sign of fluid overload
- ◆ Tense abdominal distension
- ◆ A complication that necessitates an IV infusion
- ◆ A need for feeding by NGT
- ◆ Significant refeeding diarrhoea leading to weight loss

NOTE: It is common for patients to have some change in stool frequency when their diet changes. This does not need to be treated unless there is weight loss. Having several loose stools without weight loss is not a criterion for moving back to the stabilisation phase.

Criteria to Move from the Transition Phase to Rehabilitation in the OTP

- ◆ A good appetite: Passes the appetite test and takes more than 75 percent of the daily RUTF ration.
- ◆ Oedema reduced to moderate (++) or mild (+). If wasting with bilateral pitting oedema, oedema should completely disappear.
- ◆ Medical complications are resolving.
- ◆ Clinically well and alert.

Criteria to Move from the Transition Phase to the Rehabilitation Phase in SC (for the Very Few Exceptions Who Cannot Transition to RUTF)

- ◆ A good appetite: Takes all the F-100 prescribed for the transition phase (150 kcal/kg/day).
- ◆ Oedema reduced to moderate (++) or mild (+). If wasting with bilateral pitting oedema, oedema should completely disappear.
- ◆ Medical complications are resolving.
- ◆ Clinically well and alert.

Rehabilitation Phase (Catch-up Diet for Rapid Growth)

- ◆ In the rehabilitation phase, a vigorous approach to feeding is required to achieve very high energy intakes and rapid weight gain of > 10 g/kg/day. RUTF or F-100 are used during the rehabilitation phase.
- ◆ **Patients progressing to the rehabilitation phase who consume RUTF should be transferred from SC to OTP and monitored weekly at a facility close to his/her home. Refer to section 4.1 of this guideline for case management in OTP.**
- ◆ A very small number of patients who progress from the transition phase will require to continue with the rehabilitation phase in SC.

Dietary Treatment for the Small Proportion in Need of Inpatient Rehabilitation

- ◆ Provide F-100 according to patient's body weight.
- ◆ Give feeds every four hours of F-100 per day.

Quantities of F-100 during Rehabilitation

- ◆ Give 200 ml of F-100 (200 kcal) per kg of bodyweight per day.
- ◆ Use the reference tables in [Annex 7](#) for the volume of F-100 to give per feed in the inpatient rehabilitation phase, according to patient's bodyweight.

Preparation of F-100

- ◆ Prepare F-100 feeds as indicated on [Table 18](#).

Feeding Procedure in Rehabilitation Phase

- ◆ Feed by cup and saucer.
- ◆ Breastfed children should be offered breast milk on demand before being fed F-100.
- ◆ After the feed, always offer additional F-100 if the patient takes all the feed given quickly and easily. The patient should be allowed to take as much as he/she wants.

Monitoring during Rehabilitation in the SC

Monitor and record the following parameters on the SC Multi-chart:

- ◆ Weight (daily)
- ◆ Degree of oedema: 0, +, ++, or +++ (daily)
- ◆ Body temperature, respiration, and pulse (two times per day)
- ◆ Standard clinical signs, such as stool, vomiting, dehydration, cough, and respiration (daily)
- ◆ MUAC (each week)
- ◆ Other data, such as absence, refusal of feed, etc. (daily)
- ◆ Findings of full medical examinations (daily)
- ◆ If a patient develops any signs of a medical complication while receiving treatment in rehabilitation phase, he/she should be referred to the stabilisation phase.

4.2.5 Provide Sensory Stimulation and Emotional Support

Sensory stimulation should be provided to the children throughout the period they are in SC, including:

- ◆ Tender loving care
- ◆ A cheerful, stimulating environment
- ◆ Structured play therapy for 15–30 minutes per day
- ◆ Maternal involvement, when possible, for comforting, feeding, bathing, and playing with the child

The Environment

- ◆ The room should be brightly coloured, with decorations that interest children.
- ◆ A radio can provide background music.
- ◆ The atmosphere in the room should be relaxed, cheerful, and welcoming.

Play Activities

- ◆ The mother should engage with the child by talking, singing, playing simple games, and gazing into the child's eyes with a smile. Love and affection are important for quick recovery from SAM.
- ◆ Play materials (e.g., toys) should be available for the children. These can be made locally by mothers while their children are in the SC to keep them busy and stimulate them.

Physical Activities

- ◆ Use tins as a drum and bang them with sticks.
- ◆ Put things 'in' and 'out' of a cup and teach these words whilst doing the action.
- ◆ Build towers with small blocks of wood.
- ◆ Make a ball (e.g., stuff a sock) and throw or kick the ball.
- ◆ Sing songs with actions (e.g., clapping hands).
- ◆ Play games like counting toes.
- ◆ Look at and talk about pictures.
- ◆ Teach parts of the body, or the names of clothes, when dressing.
- ◆ Teach words like 'water' and 'splash' when bathing.

See Annex 8 for examples of toys for children.

4.2.6 Failure to Respond to Treatment

Some patients in SC may fail to respond to treatment or their condition may deteriorate. The most frequent causes of failure to respond to inpatient treatment are listed in Table 19.

Table 19. Frequent Causes of Failure to Respond to Inpatient Treatment

Problems related to the health facility:

- Lack of adherence to SAM treatment protocols
- Poor environment for the patient with SAM
- Failure to complete the Multi-chart correctly resulting in data gaps relevant to the patient’s progress
- Insufficient staff (particularly at night) or inadequately trained staff
- Inadequate supervision and constant rotation of staff in the treatment facility
- Inaccurate weighing scales
- Incorrect feed preparation, or feeds given incorrectly

Problems related to the individual child:

- Infections, especially diarrhoea (amoebiasis, giardiasis, dysentery), pneumonia, TB, urinary infection, otitis media, malaria, HIV/AIDS, schistosomiasis, and/or hepatitis/cirrhosis
- Other serious underlying diseases or conditions, such as congenital abnormalities (e.g., Down’s Syndrome), neurological damage (e.g., cerebral palsy), or inborn errors of metabolism
- Insufficient feeds given
- Psychological trauma (particularly in families living with HIV and in refugee situations)
- Vitamin and mineral deficiencies
- Rumination
- Malabsorption

Patients who meet any of the criteria on [Table 20](#) should be diagnosed as failing to respond to treatment.

Table 20. Criteria for Failure to Respond to Treatment

Criteria	Time after Admission
Failure to regain appetite	Day 4
Failure to start to lose oedema	Day 4
Oedema still present	Day 10
Failure to gain at least 5 g/kg bodyweight	Day 10

- ◆ The patient with failure to respond to treatment requires extensive medical evaluation to look for causal factors; overall case management practices should also be reviewed (e.g., evaluation of adherence to treatment protocol, availability of trained staff).
- ◆ Record ‘Failure to respond to treatment’ on the individual treatment chart and schedule the patient to be seen by a senior experienced clinician or send to a referral hospital.
- ◆ Corrective measures should be taken to strengthen specific areas that need improvement in the facility’s SAM management practice while ensuring that treatment protocols are adhered to and that staff are supervised adequately.

Required Actions When Patients Fail to Respond to Treatment

- ◆ Keep accurate records of all patients who fail to respond to the treatment. These records should include, at a minimum, patient's age, sex, date of admission, MUAC on admission, principal diagnosis, treatment, and where appropriate, date, time, and apparent cause of death.
- ◆ Systematically examine the common causes of failure to respond and identify areas where case management practices should be improved to rectify problems.
- ◆ If the above actions are not immediately successful, an external evaluation should be conducted by an expert experienced in the management of SAM with medical complication. An investigation into the organisation and application of the treatment protocol should be carried out as part of the evaluation.
- ◆ Review staff supervision with refresher training, if necessary.
- ◆ Recalibrate scales.

4.2.7 Transfer to OTP and Discharge of Children 6-59 Months

Table 21. Criteria for Discharge and Transferring Children 6-59 Months

1. Stabilised and Transferred to OTP:

- ◆ Appetite has returned (passed a RUTF appetite test; the child eats more than 75% of the daily RUTF prescription) and start of weight gain.
- ◆ Medical complications are resolving.
- ◆ Bilateral pitting oedema is decreasing (if severe wasting with bilateral pitting oedema admission, bilateral pitting oedema is completely resolved).
- ◆ Clinically well and alert.

2. Discharged after Full Recovery (CURED) in SC:

Note: The same anthropometric indicator that is used to identify and confirm SAM on admission should be used to determine recovery and discharge from treatment.

If admitted with bilateral pitting oedema, discharge cured when:

- ◆ No bilateral pitting oedema for 2 consecutive visits
- ◆ **AND** MUAC \geq 12.5 cm or WFH/WFL \geq -2 z-score
- ◆ **AND** Clinically well and alert

If admitted with bilateral pitting oedema, discharge cured when:

- ◆ MUAC \geq 12.5 cm
- ◆ **AND** No bilateral pitting oedema
- ◆ **AND** Clinically well and alert

If admitted based on WFH/WFL, discharge cured when:

- ◆ WFH/WFL \geq -2 z-score
- ◆ **AND** No bilateral pitting oedema
- ◆ **AND** Clinically well and alert

Table 22. SC Exit Categories

Category	Definition
Cured	Has reached the discharge criteria for SAM treatment, i.e., the special cases that were not referred to OTP (see Table 21).
Died	Dies while receiving treatment in the SC.
Defaulted	Absent for two consecutive days. Default should be confirmed.
Non-responder	Patient who remained in treatment in the SC does not reach the SAM discharge criteria after 16 weeks (4 months) in treatment.
Stabilised	Condition has stabilised and referred to continue treatment in OTP.
Transferred-out	Moved to another facility for further medical care or moved out to receive care in another SC

4.2.8 Discharge Procedure from SC to the OTP

- ◆ Give feedback to the caregiver on the final treatment outcome.
- ◆ Ensure that the caregiver understands how to use continuing medications.
- ◆ Give a Referral Slip to the caregiver to bring to the OTP. Ensure that all medications given and important instructions to be continued in OTP are indicated.
- ◆ Update the SC Multi-chart and Registration Book for SAM with the exit category (see Table 22).
- ◆ Counsel the caregiver on the importance of continuing care in the OTP and indicate which OTP the caregiver should attend and when.
- ◆ Give adequate RUTF to last until the next OTP appointment at the site nearest to the patient's home.
- ◆ Counsel on how to use the RUTF.
- ◆ Counsel on WASH, good nutrition, and IYCF practices.
- ◆ Advise the caregiver to immediately go to the nearest health facility if they experience any signs of deterioration.

4.2.9 Treatment and Care of Infants 0-6 Months of Age

Admission Criteria and Entry Categories for Infants 0-6 Months of Age

Table 23. SC Admission Criteria for Infants 0-6 Months

- ◆ Any grade of bilateral pitting oedema (+, ++ or +++)

OR

- ◆ WFL < -3 z score

OR

- ◆ Recent weight loss or failure to gain weight.

OR

- ◆ Ineffective feeding (attachment, positioning and suckling) directly observed for 15-20, minutes, ideally in supervised separate area.

OR

- ◆ Presence of any of the following medical complications:

- ◆ Poor appetite
- ◆ Intractable vomiting
- ◆ Convulsions
- ◆ Lethargy, not alert
- ◆ Unconsciousness
- ◆ High fever
- ◆ Lower respiratory tract infection
- ◆ Dehydration
- ◆ Persistent diarrhoea
- ◆ Severe anaemia
- ◆ Hypoglycaemia
- ◆ Hypothermia
- ◆ Severe skin lesions
- ◆ Eye signs of vitamin A deficiency

OR

- ◆ Any medical or social issue needing more detailed assessment or intensive support (e.g. disability, depression of caregiver, or other adverse social circumstances).

Table 24. SC Entry Categories

Category	Definition
New admission	New case who meets the admission criteria for SC (see Table 23).
Readmission	Re-admitted for treatment in SC due to relapse or a returned defaulter who meets the admission criteria.
Relapse	Cured within the past 3 months and now meets the admission criteria for SC (see Table 23).
Returned defaulters	Defaulted within the past 3 weeks and has returned to continue treatment in SC. A returned defaulter should be re-admitted if they meet the admission criteria.
Transfers-in	Has moved in from another facility where s/he was receiving care in SC.

Management of the Breastfed Infant 0-6 Months

- ◆ Prioritize re-establishing effective exclusive breastfeeding.
- ◆ Promote and support the mother or caregiver to breastfeed.
- ◆ Assess the physical and mental health status of mothers or caregivers and provide relevant treatment and support.

Routine Medicines and Supplements

Choice of Broad-Spectrum Antibiotics

- ◆ The choice of broad-spectrum antibiotics for infants 0-6 months is the same as that of children 6-59 months. See Annex 3.

Ferrous Sulphate

- ◆ Provide daily doses of iron syrup orally (see dosage instructions on Table 25).

Table 25. Doses of Iron Syrup if F-100 Is Used in Rehabilitation

Weight of Child	Dose of Iron Syrup: Ferrous Fumarate 100mg per 5ml (20mg elemental iron per ml)
3–6 kg	0.5 ml
6–10 kg	0.75 ml
10–15 kg	1 ml

Note that the above dosages are very small (less than 1/4 teaspoon) and need to be measured with a syringe.

Dietary Treatment

The objective is to supplement the child's breastfeeding with therapeutic milk while stimulating breast milk production.

- ◆ The infant should be breastfed as frequently as possible. Breastfeed every 3 hours for at least 20 minutes (more if the child cries or demands more).
- ◆ Between 30 minutes and 1 hour after a normal breastfeeding session, give maintenance amounts of therapeutic milk.
 - ◆ Give F-100-Diluted to breastfed infants without bilateral pitting oedema. F-100-Diluted is prepared by adding 30% water to dilute full strength F-100. [See Table 25](#).
 - ◆ Give F-75 to infants with bilateral pitting oedema and change to F-100-Diluted when the oedema is resolved.
 - ◆ Never give full strength F-100 to infants 0-6 months old.
 - ◆ Quantities of F-100-Diluted or F-75 to Give the Breastfed Infant
 - ◆ Give F-100-Diluted or F-75 every 3 hours (8 feeds per day).
 - ◆ Use the reference tables [\(see Annex 9\)](#) for amounts of F-100-Diluted or F-75 to give to infant.
 - ◆ Use the supplementary suckling technique [\(see Figure 5\)](#). The quantity of F-100-Diluted or F-75 should not be increased as the child starts to gain weight.

Preparation of F-100-Diluted using the New Tin Formulation

1. Boil water to make it safe for drinking.
2. Ensure that the water temperature is not below 70°C (i.e., cooled for not less than 3 – 5 minutes after boiling).
3. Add water to F-100 therapeutic milk. [See Table 26](#).
4. Whisk the mixture vigorously until the powder dissolves in the water.
5. Cool the prepared milk to feeding temperature before administering.
6. Give the feed based on the child's body weight. [See Annex 9](#).

Table 26. Preparation of F-100-Diluted Feeds using the New Tin Packaging

Blue Scoops of F-100	Water (ml)	*Volume of F-100-Diluted Milk (ml)
1	33	38
2	65	75
3	98	113
4	130	152
5	163	189
6	195	226
7	228	264
8	260	304
9	293	339
10	325	377
20	650	754
Tins of F-100	Water (ml)	*Volume of F-100-Diluted Milk (ml)
1 Tin (400 grams)	2860	2805
2 Tins (800 grams)	5720	5611

*Note that the F-100 Diluted milk volumes provided are estimates; the health care provider should measure the amount of milk to feed the child based on his/her body weight.

Supplementary Suckling Technique

Use the supplementary suckling technique to re-establish or commence breastfeeding, and to provide maintenance amounts of F-100-Diluted to severely malnourished infants. The technique entails having the infant suckle the breast while also taking supplementary F-100-Diluted through a fine tube that runs alongside the nipple. The infant is nourished by the supplementary F-100-Diluted and the suckling stimulates the breast to produce more milk. Follow the steps below:

- ◆ The caregiver holds a cup of F-100-Diluted or F-75.
- ◆ One end of an NGT (size no 8) is put in the cup and the other end of the tube is placed on the breast, at the nipple.
- ◆ The infant is offered the breast, ensuring proper attachment.
- ◆ The cup should be held 5–10 cm below the level of the nipple for easy suckling. If the child has a strong suckle, the cup can be lowered to up to 30 cm below nipple-level.

Figure 5. Supplementary Suckling Technique



After feeding is completed, the tube is flushed through with clean water using a syringe. It is then spun (twirled) rapidly to remove the water in the lumen of the tube by centrifugal force. If possible, the tube is then exposed to direct sunlight to kill bacteria.

Individual Monitoring

Monitor and record the following parameters on the SC Multi-chart:

- ◆ Weigh the infant 0-6 months with a scale graduated to within 10 g (or 20 g)
- ◆ Degree of oedema (0, +, ++, or +++)
- ◆ Body temperature, pulse, and respiration
- ◆ Standard clinical signs: stool, vomiting, dehydration, cough, and respiration.
- ◆ Other relevant data, e.g., if the child is absent, vomits or refuses a feed; whether the patient is fed by NGT or given an IV infusion or transfusion.

Supportive Care for Mothers

- ◆ Provide mothers with supportive care for breastfeeding.
- ◆ Focus on creating conditions that will facilitate and increase breastfeeding, such as establishing safe “breastfeeding corners” for mothers and infants, one-on-one counselling, and mother-to-mother support.
- ◆ Assess the mother’s mental health for signs and symptoms of depression, anxiety, trauma, or other emotional distress. Traumatized and depressed mothers may have difficulty caring for their infants and require mental and emotional support to effectively breastfeed.
- ◆ Assess the mother’s nutritional status using MUAC, if < 23.0 cm, admit in the TSFP (see “Management of MAM” on chapter 5)
- ◆ Explain to the mother the different treatment steps her child will go through. Try to strengthen the mother’s confidence and discourage self-criticism for her perceived inability to provide adequate breast milk.
- ◆ Alert the mother of the risk of a new pregnancy during this period.

Adequate Nutrition and Supplementation for Breastfeeding Mothers

Breastfeeding mothers need about 500 kcal per day of extra energy. Essential micronutrients in breast milk come from the mother’s diet or micronutrient supplements. It is important that the mother’s nutrient and energy needs be met. The mother should consume at least 2,500 kcal per day. The health facility should provide nutritious food for the mother. Dehydration may interfere with breast milk production; it is important to ensure that the mother drinks at least 2 litres of water per day.

Discharge and Exit Categories for the Breastfed Infant 0-6 Months

Table 27. Discharge Criteria for the Breastfed Infants 0-6 Months

Discharge when the following criteria has been met:

- Successful re-lactation and effective breastfeeding has been achieved.
- Gaining weight on exclusive breastfeeding (i.e., more than 5 g/kg/day for at least 3 successive days).
- No bilateral pitting oedema.
- Clinically well and alert.
- Infant has been checked for immunization and other routine interventions.
- Mothers or caregivers have been linked with community-based follow-up and support.

Table 28. SC Exit Categories

Category	Definition
Cured	Has reached the discharge criteria for SAM treatment (see Tables 27 for breastfed and Table 29 non-breastfed infants).
Died	Dies while receiving treatment in the SC.
Defaulted	Absent for two consecutive days. Default should be confirmed.
Non-responder	Does not reach the SAM discharge criteria after 16 weeks (4 months) in treatment.
Transferred-out	Moved to another facility for further medical care, or, moved out to receive care in another SC.

Discharge Procedure

- ◆ Give feedback to the caregiver on the final treatment outcome of the infant.
- ◆ Ensure that the caregiver understands how to use and continue medications.
- ◆ Measure weight and assess for bilateral pitting oedema in the infant.
- ◆ Be sure to update the SC Multi-chart and Registration Book for SAM with the treatment outcome. See Table 28.
- ◆ If the mother has a MUAC of < 23.0 cm, provide a 2-week ration of the specialised nutritious food, complete a Referral Slip and refer her to continue with TSFP at the nearby health facility.
- ◆ Counsel on the importance of continuing care in the TSFP and indicate which TSFP site the caregiver should attend and when.

Management of the Non-Breastfed Infant 0-6 Months

- ◆ Prioritize establishing, or re-establishing effective exclusive breastfeeding by the mother. If available and culturally accepted, support the female caregiver with re-lactating.
- ◆ If there is no realistic prospect of the infant being breastfed, manage the infant using F-75 (if bilateral pitting oedema) or F-100-Diluted.

Treatment of non-breastfed infants 0-6 months is divided into stabilisation phase, transition, and rehabilitation phase.

STABILISATION PHASE

Routine Medicines and Supplements

Choice of Broad-Spectrum Antibiotics

The choice of broad-spectrum antibiotics for infants 0-6 months is the same as that of children 6-59 months. [See Annex 3.](#)

Ferrous Sulphate

Provide daily doses of iron syrup orally ([see Table 25](#)).

Dietary Treatment

The dietary treatment for non-breastfed infants is the same as that of breastfed infants. [See “Dietary Treatment” on page 76.](#)

Feeding Procedure

Apply the same feeding procedure for infants 0-6 months as that of children 6-59 months in the stabilisation phase ([see “Feeding Procedure in Stabilisation Phase” on page 62](#)).

Individual Monitoring

Monitor and record the following parameters on the SC Multi-chart:

- ◆ Weigh the infant 0-6 months with a scale graduated to within 10 g (or 20 g)
- ◆ Degree of oedema (0, +, ++, or +++)
- ◆ Body temperature, pulse, and respiration
- ◆ Standard clinical signs: stool, vomiting, dehydration, cough, and respiration.
- ◆ Other relevant data (e.g., if the child is absent, vomits, or refuses a feed; whether the patient is fed by an NGT or is given IV infusion or transfusion)

Criteria to Progress from the Stabilisation Phase to the Transition Phase

Two criteria must be met for children to progress from the stabilisation to the transition phase:

- ◆ Return of appetite.
- ◆ Lessening of oedema, which normally consists of appropriate and proportionate weight loss as oedema starts to subside (children with severe oedema (+++) should remain in the stabilisation phase until their oedema can be categorized as moderate (++)).

TRANSITION

Routine Medicines and Supplements

Routine antibiotic therapy should be continued during the transition phase until the child is transferred to the rehabilitation phase.

Dietary Treatment

- ◆ Use the standard protocol for older children in the transition phase, with one modification—use only F-100-Diluted.
- ◆ Increase the volume of F-100-Diluted feeds by 30%, as compared to the amount given during the stabilisation phase. [See Annex 9](#) for F-100-Diluted to give to non-breastfed infants in the transition phase.

Individual Monitoring

Continue with monitoring during transition as outlined for children 6-59 months. [\(see “Monitoring during Transition” on page 66\).](#)

- ◆ Criteria to Progress from the Transition Phase to the Rehabilitation Phase
- ◆ The following criteria should be met before the infant progresses from the transition to the rehabilitation phase:
- ◆ A good appetite, defined as taking almost all (at least 90%) of the F-100-Diluted prescribed for the transition phase.
- ◆ Complete loss of bilateral pitting oedema.
- ◆ Completed a minimum 2-day stay in the transition phase.
- ◆ No medical complications.

REHABILITATION PHASE

Dietary Treatment

Use the standard protocol for older children in the rehabilitation phase, with the following modifications:

- ◆ Use F-100-Diluted.
- ◆ Infants should receive twice the volume of F-100-Diluted per feed, as compare to the amount given during the stabilisation phase. [See Annex 9](#) for the amounts of F-100-Diluted to give to non-breastfed infants in the rehabilitation phase.

Individual Monitoring

Continue with monitoring in rehabilitation phase as outlined for children 6-59 months. [\(see “Monitoring during Rehabilitation in the SC” on page 68\).](#)

Discharge Criteria and Exit Categories for the Non-Breastfed Infant 0-6 Months

Table 29. Discharge Criteria for the Non-Breastfed Infants 0-6 Months

Discharge when the following criteria has been met:

- Infant is feeding well with the replacement feed.
- Has adequate weight gain and has a WFL ≥ -2 z-score.
- No bilateral pitting oedema.
- Clinically well and alert.
- Infant has been checked for immunization and other routine interventions.
- Mothers or caregivers have been linked with community-based follow-up and support

Discharge Procedure

Follow the same discharge procedure as that of breastfed infants. [See “Discharge Procedure” on page 79](#)

5 Management of Moderate Acute Malnutrition (MAM)

This chapter describes two types of Supplementary Feeding Programmes (SFPs) used to manage MAM. The chapter highlights on the principles, criteria, and process of implementation. It is recommended that MAM interventions are linked with SAM treatment, IYCF, and social behaviour change communication (SBCC), WASH and other health and nutrition interventions. Households of MAM patients should also be referred for PSNP.

5.1 Principles of SFPs

- ◆ SFPs aim to prevent and rehabilitate acute malnutrition.
- ◆ A take-home specialised nutritious food is provided to the patient, with follow-up visits conducted at a nearby health facility every two weeks.
- ◆ For SFPs to achieve the intended outcomes, it is critical that effective and appropriate linkages are made with food security interventions to avoid the risk of sharing the specialised nutritious food with other members of the household.
- ◆ **Table 30** below shows the two types of SFPs, Blanket Supplementary Feeding Programme (BSFP) and Targeted Supplementary Feeding Programme (TSFP).

Table 30. Types of SFPs

BSFP	TSFP
<ul style="list-style-type: none"> ■ A supplementary ration is provided to everyone in an identified vulnerable group for a defined period. For example, all children 6-23 or 6-59 months, or all pregnant and lactating women, regardless of their nutritional status. ■ Anthropometric indicators are <u>not</u> used to determine those admitted or discharged from treatment. However, beneficiaries are screened to identify those in need of SAM treatment. 	<ul style="list-style-type: none"> ■ A supplementary ration is targeted to individuals with MAM in specific vulnerable groups. The vulnerable groups usually include children 6-59 months and pregnant and lactating women with infants under 6 months of age who have MAM. ■ TSFP might also include individuals with special needs such as people living with HIV (PLHIV), tuberculosis (TB) patients and the older people with MAM. ■ Specific anthropometric criteria are used to determine those admitted and discharge from treatment.

5.2 Blanket Supplementary Feeding Programme (BSFP)

When Should BSFP be Established?

BSFP is implemented when the Global Acute Malnutrition (GAM) levels are high and food insecurity is severe. It can be implemented as part of an emergency response or as a response to seasonal shocks to prevent spikes in acute malnutrition. BSFP aims to prevent acute malnutrition or further deterioration of nutritional status among identified vulnerable groups. See Annex 10 for the decision tree on MAM.

Admission Criteria and Discharge Criteria

- ◆ The admission and discharge criteria for BSFP does not rely on anthropometric indicators.
- ◆ Define the targeted groups, for example, all children aged 6-23 months or 6-59 months, or all PLW.
- ◆ BSFP normally operates for a period of about 3-6 months, after the situation has stabilized, it is closed and those registered are discharged.

Treatment and Follow-Up Procedure in BSFP

- ◆ Register all individuals within the defined target group for the BSFP.

Screen the individuals by measuring MUAC and assessing for bilateral pitting oedema. The screening is conducted to identify cases who may require referral for SAM management. It is not used for admission to the BSFP.

- ◆ Provide a two-week supply of specialised nutritious foods to all those registered for BSFP.
- ◆ Monitor individual nutritional status by measuring MUAC and assessing for bilateral pitting oedema during each bi-weekly follow-up visit.
- ◆ Ensure provision of other health services such as: vitamin A supplementation, deworming, insecticide treated nets, water purification tablets, and health and nutrition education.

When Should BSFP be Closed?

BSFP should be closed when GAM is below 10% with no aggravating factors and the following met:

- ◆ The general food distribution is reliable and adequate, or food security is acceptable.
- ◆ Effective public health and disease control measures are in place.
- ◆ No seasonal deterioration of nutritional status is expected.
- ◆ The population size is stable, with no new displacement expected.
- ◆ There is availability and access to WASH facilities.
- ◆ Nutrition sensitive interventions are in place.

5.3 Targeted Supplementary Feeding Programme (TSFP)

TSFP should be integrated with routine health services and OTP services where possible. It aims to: 1) treat patients with MAM; 2) prevent deterioration from MAM to SAM; 3) prevent mortality associated with MAM; and 4) prevent deterioration in maternal nutritional status and subsequently prevent poor birth outcomes.

5.3.1 Admission Criteria and Entry Categories

Admission Criteria

Table 31. TSFP Admission Criteria

Children 6–59 Months	Pregnant Women and Lactating Women (up to 6 Months Postpartum)
<ul style="list-style-type: none"> ■ MUAC \geq11.5 to $<$12.5 cm ■ WFH \geq -3 to $<$-2 Z scores ■ No bilateral pitting oedema <p>AND</p> <ul style="list-style-type: none"> ■ No medical complications ■ Clinically well and alert 	<ul style="list-style-type: none"> ■ MUAC $<$ 23.0 cm

Entry Categories

Table 32. TSFP Entry Categories

Category	Definition
New admissions	New case who meet admission criteria for TSFP (see Table 31).
Readmission	Re-admitted for treatment in TSFP due to relapse or a returned defaulter who meets the admission criteria.
Relapse	Cured within the past 3 months and now meets the admission criteria for TSFP (see Table 31).
Returned defaulters	Defaulted within the past 3 months and has returned to continue treatment in TSFP. A returned defaulter should be re-admitted if they meet the admission criteria.
Transfers-in	Has moved in from another facility where s/he was receiving TSFP.

5.3.2 Admission Procedure

Once a decision has been made to admit the patient in TSFP, the steps below should be completed during the admission process.

Step 1: Take Anthropometric Measurements and Confirm Nutritional Status

- ◆ Determine the age of the patient based on the health card or caregiver's recall.
- ◆ Check for bilateral pitting oedema and record the findings.
- ◆ Measure MUAC of the patient and record the measurement.
- ◆ Measure and record weight of the patient.
- ◆ Measure and record height or length (if at HC level).
- ◆ Determine WFH/WFL z-score using the reference charts (if at HC level).
- ◆ Classify nutritional status.

Step 2: Admit to the TSFP

- ◆ Explain the TSFP treatment procedures and care that will be provided.
- ◆ Decide which TSFP admission entry category to assign to the patient (see Table 32).
- ◆ Assign a registration number if the patient is a new admission.
- ◆ Give routine medicines in TSFP. See Table 33 and Table 34.
- ◆ If scheduled immunisations have not been given, administer them.
- ◆ Give the specialised nutritious food (see Table 35 for Fortified Blended Food (FBF) and Table 36 for Ready-to-Use Supplementary Food (RUSF)
- ◆ Complete the TSFP Treatment and Follow-up Card. See Annexes 19 and 20.
- ◆ Register the patient in the registration book for MAM treatment. Be sure to use the appropriate registration book for children or PLW.
- ◆ Provide nutrition counselling and education, including IYCF.
- ◆ Issue the patient with the TSFP Identification Card. See Annex 21.
- ◆ Advise the caregiver or patient when to return for the next visit.
- ◆ Refer to other health and nutrition services such as antenatal care, post-natal care, and PSNP.

5.3.3 Medical Treatment and Care

- ◆ Check and provide routine medications such as vitamin A, deworming, and routine immunizations. See [Table 33](#) and [Table 34](#) for the routine medications to give to children and PLW in TSFP.

Table 33. Routine Medication and Treatment for Children in TSFP

Name of Medicine	When	Age	Dose	Prescription
Albendazole Or	On admission	24-59 months	400 mg	Single dose on admission
Mebendazole	On admission	24-59 months	500 mg	Single dose on admission
Measles vaccination	On admission if the child has not received the measles vaccine before.	9-59 months	Refer to the national EPI guidelines	Refer to the national EPI guidelines
Vitamin A	On admission	6-59 months	6-11 months: 100,000 IU 12-59 months: 200,000 IU	Single dose, if not provided within the last six months

Table 34. Routine Medication and Treatment for PLW in TSFP

Name of Medicine	When	Physiological Status	Dose	Prescription
Albendazole Or	Second or Third trimester	Pregnant or lactating	400 mg	Single dose on admission, if not already received in antenatal or postnatal care.
Mebendazole	Second or Third trimester	Pregnant or lactating	500 mg	Single dose on admission, if not already received in antenatal or postnatal care.

5.3.4 Nutrition Rehabilitation

- ◆ There are a range of specialised nutritious foods used for nutrition rehabilitation of patients with MAM. See [Annex 11](#).
- ◆ Fortified blended food (FBF) e.g., Super Cereal Plus, and ready-to-use supplementary food (RUSF) are the two types of specialised nutritious foods used for management of MAM in Ethiopia.
- ◆ Ideally RUSF is used for nutrition rehabilitation of children with MAM, if not available, FBF should be used.

Fortified Blended Food (FBF)

- ◆ FBF requires to be cooked with clean drinking water prior to consumption.
- ◆ Give a take-home ration of FBF to the patient with MAM. See Table 35 for the amount to give.

Table 35. FBF Ration Size per Patient

Target Group	Daily Quantity	Ration for Two Weeks	Ration for one Month
Children 6-59 months	200 g	3.0 kg	6.0 kg
PLW	250 g	3.75 kg	7.5 kg

STEP 1: Health, nutrition, and hygiene education

- ◆ Conduct health, nutrition, and hygiene education.
- ◆ Explain the following key messages on FBF to the caregiver or patient:
 - ◇ The ration is for the patient with MAM. Do not share with the rest of the family.
 - ◇ Do not give FBF to infants 0-6 months. They should be exclusively breastfed.
- ◆ If the child is 6-23 months, continue to breastfeed on demand. Mother's milk is best for infants and young children.
 - ◇ FBF should be consumed in addition to, and not as a substitute for, the household diet and breastfeeding.
 - ◇ Ensure that the FBF porridge is consumed within 30 minutes of cooking, this will avoid contamination.
 - ◇ Once the FBF has been prepared, the remainder should be store in a closed bag in a cool, dry and hygienic place, away from insects or rodents.

STEP 2: Demonstrate how to prepare the FBF porridge

- ◆ Demonstrate to the caregivers or patient how to prepare FBF at home, see Figure 6 on the preparation of the FBF porridge.

STEP 3: Distribute the FBF

- ◆ Distribute the ration of FBF to the caregiver or patient, see Tables 34 and 35 for the amount to give.


Figure 6. Preparation of the FBF Porridge

- ◆ Wash hands thoroughly using water and soap.

Step 1:



Step 2:




Step 3:



Step 4:



- ◆ Also wash the utensils with soap and water.
- ◆ If the water to be used for preparation is not clean safe drinking water, it should be boiled for 15-20 minutes before adding the FBF.
- ◆ Open the FBF bag carefully.
- ◆ Mix 1 part (e.g., cup) of the FBF with 4 parts (e.g., cups) of water.
- ◆ Cook mixture over low heat for 5-10 minutes stirring constantly.
- ◆ Prepare to a thick porridge (not too thick or too thin).
- ◆ Serve warm and consume within 30 minutes.
- ◆ Tightly close the bag of remaining FBF.



Good thickness

Poor thickness

RUSF

- ◆ RUSF is a ready-to-use food that does not require to be prepared, cooked, or mixed with water prior to consumption.
- ◆ See Table 36 for RUSF to give.

Table 36. RUSF Ration Size per Patient

Food Item	Daily Quantity	For Two Weeks	For One Month
RUSF	1 Sachet	15 Sachets	30 Sachets

Health, nutrition, and hygiene education

- ◆ Explain the following to the caregiver or patient:
 - ◆ Gently mix by pressing the sachet for 30 seconds.
 - ◆ Make a small opening in the corner of the sachet. The patient can eat directly from the sachet.
 - ◆ Drink clean safe drinking water while eating the RUSF to keep hydrated.
 - ◆ If the child is breastfeeding, continue to offer breast milk on demand.
 - ◆ RUSF is complementary to the normal meals not a replacement, therefore should be consumed between normal meals, continue consuming other nutritious foods during the treatment.
 - ◆ RUSF is ready to be used, therefore should not be mixed with other foods.
 - ◆ If the patient has diarrhoea, do not stop feeding. Continue to breastfeed and give extra food and clean water.
 - ◆ Once opened, store the RUSF sachet in a clean, cool and dry place

5.3.5 Monitoring During TSFP Follow-Up Visits

- ◆ Monitor the following parameters and record information on the TSFP Treatment and Follow-up Card at each follow-up visits.
 - ◆ MUAC
 - ◆ Weight
 - ◆ Height/length (if at HC level, measure once a month)
 - ◆ WFH/WFL z-score (if at HC level, check on admission, every month, and on discharge)
 - ◆ Presence of bilateral pitting oedema

-
- ◆ Weight gain (patients who lose weight, do not gain weight, or have weight fluctuations should receive special attention during home visits)
 - ◆ Inform the caregiver or patient on the progress of treatment.
 - ◆ Conduct individual and/or group counselling on health, WASH, and IYCF.
 - ◆ Indicate any necessary follow-up actions on the TSFP Treatment and Follow-up Card, such as:
 - ◆ Home visits for patients requiring special attention
 - ◆ Transfer to OTP for patient whose nutritional status deteriorates to SAM
 - ◆ Prioritize patients for a home visit if the patient:
 - ◆ Is losing or not gaining weight
 - ◆ Is absent or defaulted treatment (defaulters are patients absent for 2 consecutive visits).

Failure to Respond to Treatment

HIV/AIDS or TB may be the underlying cause of acute malnutrition, it is recommended that patients who are failing to respond to MAM treatment are offered HIV counselling and testing, and TB testing. Other reasons for failure to respond to treatment may include:

- ◆ Problems with the application of the protocol
- ◆ Home and social circumstances of the individual
- ◆ Excessive sharing of the TSFP ration

Steps to take if a patient with MAM is failing to respond to treatment:

1. Review the patient's TSFP Treatment and Follow-up Card to confirm that the TSFP protocols are adhered to correctly. Ensure that the details on the card are correct.
2. Conduct a full medical examination and nutrition assessment. If the MAM patient presents with any signs of medical complications or nutrition deficiencies, conduct further investigation and/or provide treatment.
3. Conduct a home visit to assess the patient's home and social situation. Assess if the patient is taking the treatment correctly, if the treatment is acceptable and tolerated, or if it is being shared with other members of the household. [See Annex 1](#) for home visit checklist.

5.3.6 Discharge Criteria and Exit Categories

Table 37. TSFP Discharge Criteria

Children 6–59 Months	Pregnant Women and Lactating Women (up to 6 Months Postpartum)
<p>If admitted based on MUAC, discharge cured when:</p> <ul style="list-style-type: none"> ◆ MUAC \geq 12.5 cm for two consecutive visits ◆ AND Clinically well and alert <p>If admitted based on WFH/WFL discharge cured when:</p> <ul style="list-style-type: none"> ◆ WFH/WFL \geq -2 z-score for two consecutive visits ◆ AND Clinically well and alert 	<p>MUAC \geq 23.0 cm for two consecutive visits</p> <p>OR</p> <p>Lactating mother whose infants reaches 6 months of age.</p>

Table 38. TSFP Exit Categories

Category	Definition
Cured	Has reached the discharge criteria for MAM treatment (see Table 37).
Died	Dies while receiving treatment in TSFP.
Defaulted	Absent for two consecutive visits. Default should be confirmed.
Non-responder	Does not reach the MAM discharge criteria after 16 weeks (4 months) in treatment.
Transferred-out	Condition has deteriorated to SAM and referred for OTP, or, moved out to receive TSFP in another facility.
Other	Pregnant woman who has delivered or a lactating woman whose infant is 6 months.

5.3.7

- ◆ Give feedback to the caregiver or patient on the final treatment outcome.
- ◆ Give the final 2-week supply of the specialised nutritious food (FBF or RUSF).
- ◆ Record the exit category (see Table 38) in the Registration Book for MAM and the TSFP Treatment and Follow-up Card.
- ◆ Counsel on WASH, good nutrition, and IYCF practices.
- ◆ Advise the caregiver or patient to immediately go to the nearest health facility if they experience any signs of deterioration.
- ◆ Refer the caregiver to PSNP and other complementary health and nutrition services.


6 Managing Acute Malnutrition in Other Vulnerable Groups

This chapter describes the management of acute malnutrition in other vulnerable groups including in the context of HIV/AIDS, TB, older people, and clients with disabilities. It also highlights the key considerations and procedures for case management.

6.1 Acute Malnutrition and HIV/AIDS

The link between nutrition and infections such as HIV/AIDS and TB is well established. Poor nutrition impairs the immune response and makes it more difficult for the body to fight infections. Infections can alter the way the body absorbs and uses nutrients, increase energy and nutrient needs, and increase nutrient losses, all of which can lead to undernutrition. See Figure 7 below

Figure 7. Cycle of Poor Nutrition and Infection



Adapted from: Regional Centre for Quality of Health Care and FANTA. 2003. Handbook: Developing and Applying National Guidelines on Nutrition and HIV/AIDS. Kampala and Washington, DC: RCQHC and FANTA/FHI 360.

.....

In People Living with HIV (PLHIV), optimal nutrition is critical to effective HIV care and treatment; when the body's immune system is compromised due to HIV or AIDS, this can contribute to malnutrition and susceptibility to infection. On the other hand, a person who is well-nourished is more likely to maintain a stronger immune system that helps fight the HIV infection. [See Annex 12](#) for the nutrient requirements for PLHIV.

The following recommendations are key when managing SAM and MAM in the context of HIV/AIDS:

- ◆ All patients failing to respond to SAM or MAM treatment should be offered HIV counselling and testing.
- ◆ Refer HIV-positive patients with SAM and MAM for treatment according to National Guidelines for Comprehensive HIV Prevention, Treatment and Care
- ◆ HIV-positive children with SAM or MAM should be given the same ART treatment regimen, in the same doses, as patients with HIV who do not have SAM or MAM.
- ◆ Children with SAM or MAM who are HIV-positive should be managed using the same specialised nutritious foods (i.e., F-75, F-100, RUTF, RUSF, or FBF) as those with SAM or MAM who are HIV-negative.
- ◆ Children with SAM or MAM who are HIV-positive should follow the same guidance for micronutrient supplementation including Vitamin A, iron, folic acid, and zinc for diarrhoea management as those with SAM and MAM who are HIV-negative.

Additionally, the following specific recommendations will apply to PLHIV who have SAM:

- ◆ Children with SAM who are HIV-positive and qualify for antiretroviral therapy (ART) should be started on treatment as soon as possible after stabilisation of medical complications.
- ◆ HIV-positive children with SAM who are started on ART treatment should be monitored closely 6–8 weeks following initiation of ART.
- ◆ HIV-positive children with SAM in whom persistent diarrhoea does not resolve with standard management should be investigated to exclude carbohydrate intolerance and infective causes, which may require different management such as modification of fluid and feed intake, or antibiotics.

6.2 Acute Malnutrition and TB

Acute malnutrition increases the risk of TB and vice-versa and, as a result, acute malnutrition is highly prevalent among people with TB ([see Figure 7](#)). Children and pregnant women are particularly vulnerable to the effects of acute malnutrition and TB. Women with TB may be at higher risk for pre-eclampsia and other complications during pregnancy. TB also increases the risk of premature birth, low birth weight, and perinatal death.

- ◆ The following should therefore be considered when managing acute malnutrition in the context of TB:
- ◆ Actively screen for acute malnutrition in TB patients and provide treatment and care for SAM or MAM based on their nutritional status.
- ◆ All patients failing to respond to SAM or MAM treatment should be tested for TB as a possible underlying cause of acute malnutrition.

- ◆ Children with SAM or MAM and TB infection should be managed using the same specialised nutritious foods (i.e., F-75, F-100, RUTF, RUSF, or FBF) as those with SAM or MAM who do not have TB infection.
- ◆ If TB is strongly suspected in a child with acute malnutrition (i.e., if the child has had contact with an adult TB patient, poor growth despite good intake, chronic cough, or a chest infection that has not responded to antibiotics):
 - ◆ Perform a Mantoux test (NOTE: false negatives are frequent).
 - ◆ Take a chest x-ray, if possible.
 - ◆ If the test is positive or there is a strong suspicion of TB, treat according to National TB Guidelines.

6.3 Acute Malnutrition in Older People (≥ 60 Years)

6.3.1 Community Outreach

- ◆ Older people include all men and women ≥ 60 years of age.
- ◆ Screen for acute malnutrition in the community by measuring MUAC and assessing for bilateral pitting oedema.
- ◆ Older people with a MUAC < 21.0 cm and/or bilateral pitting oedema should be referred for management at a nearby HP.

6.3.2 Admission and Discharge Criteria for the Management of SAM

Table 39. Admission and Discharge Criteria for SAM in Older People

Admission Criteria	Discharge Criteria
<ul style="list-style-type: none"> ■ Bilateral pitting oedema <p>OR</p> <ul style="list-style-type: none"> ■ MUAC < 18.5 cm <p>AND</p> <ul style="list-style-type: none"> ■ Appetite passed ■ Clinically well and alert 	<ul style="list-style-type: none"> ■ MUAC ≥ 18.5 cm for two consecutive visits <p>AND</p> <ul style="list-style-type: none"> ■ No bilateral pitting oedema <p>AND</p> <ul style="list-style-type: none"> ■ Clinically well and alert

6.3.3 Considerations for the Management of SAM

- ◆ Follow the SAM treatment protocol for children 6-59 months to manage SAM in older people ≥ 60 years who meet the admission criteria in [Table 39](#).
- ◆ Discharge patient to TSFP when they have a MUAC ≥ 18.5 cm and no bilateral pitting oedema.
- ◆ Report data using the “other” category of the Monthly Statistics Report for Acute Malnutrition. (See [Annex 26](#)).

6.3.4 Admission and Discharge Criteria for the Management of MAM

Table 40. Admission and Discharge Criteria for MAM in Older People

Admission Criteria	Discharge Criteria
<ul style="list-style-type: none"> ■ MUAC ≥ 18.5 to < 21.0 cm <p>AND</p> <ul style="list-style-type: none"> ■ No bilateral pitting oedema ■ Appetite passed ■ Clinically well and alert <p>OR Discharged from OTP</p>	<ul style="list-style-type: none"> ■ MUAC ≥ 21.0 cm for two consecutive visits <p>AND</p> <ul style="list-style-type: none"> ■ Clinically well and alert

6.3.5 Considerations for the Management of MAM


- ◆ Follow the TSFP treatment protocol for children 6-59 months to manage MAM in older people ≥ 60 years who meet the MAM admission criteria in [Table 40](#).
- ◆ Provide the following routine medications to older people:
 - ◆ Single high dose of vitamin A (200,000 IU) on admission to TSFP.
 - ◆ Single dose of Mebendazole or Albendazole (500 mg) on admission to TSFP.
 - ◆ Folic acid 10-20 mg for 30 days.
- ◆ Report data using the “other” age category of the Monthly Statistics Report for Acute Malnutrition.

6.4 Acute Malnutrition and Disability

- ◆ As nutrition outcomes improve and patients who previously would have died now survive, disability is becoming a more common underlying factor of admissions in acute malnutrition services. There are important links and interactions between disability and malnutrition, each can cause or contribute to the other ([see Figure 8](#)).
- ◆ The principles, criteria, and process for the management of SAM and MAM in infants 0-6 months and children 6-59 months without disabilities will apply in the care and treatment of those with disabilities.

- ◆ People with disability may not be able to reach or access food as well as people without disabilities. They may also have impaired swallowing or lose nutrients through vomiting. Cerebral palsy and cleft lip/palate are some of the common disabilities in patients with acute malnutrition. The following considerations should be made in the context of disabilities:
- ◆ Proactively screen for acute malnutrition to identify and manage disability and associated conditions for the following reasons:
 - ◆ To counsel and advise the patient or caregiver about the disability
 - ◆ To offer disability-specific feeding advice and treatment
 - ◆ To have more realistic outcome expectations (e.g., the patient may be much slower to gain weight)
 - ◆ To ensure that the patient is referred to appropriate support services/organisations
- ◆ Conduct regular home visits to households of people with disability to support and follow-up those receiving care and treatment.

Figure 8. Interactions Between Malnutrition and Disability



Source: Kerac M. et al. 2014. The interaction of malnutrition and neurologic disability in Africa. *Seminars in Pediatric Neurology*, 21 (1). pp. 42-9. ISSN 1071-9091 DOI.

7 Supply Management

This chapter describes the essential supplies for acute malnutrition and their management. It also highlights on the processes and procedures that need to be undertaken to ensure consistent availability of SAM and MAM supplies at the facility level.

7.1 Essential Supplies for Acute Malnutrition

An efficient and uninterrupted supply management should always be maintained. [Annex 13 and 14](#) provides a list the essential supplies needed for the management of SAM and MAM.

7.2 Procurement

Procurement of the essential SAM and MAM supplies should follow national and international standards and regulations.

7.3 Requesting Supplies

The quantity of supplies needed will depend on the SAM or MAM caseload, the amount normally used in each period (e.g., each quarter), frequency of requests, and the existing storage capacity.

7.3.1 Quantifying the Need

The first step in requesting supplies is to quantify the need. When quantifying needs at the health facility, woreda, zonal or regional level, the service provider should do the following:

1. Identify the item or product needed.
2. Determine the number of SAM or MAM cases receiving treatment in the area. To do this, use the routine programme data, or in the absence of routine programme data, use the most recent prevalence of SAM or MAM in the area
3. Determine the average consumption of the item or product.
4. Decide the period for which supplies are needed (i.e, one month, three months, or one year). This will be based on the storage space available, the distance to the collection point, and product expiry date.
5. Calculate the need using the caseload, average consumption, and period for which supplies are needed. Add a 10% buffer stock to the need. Consider recent events such as influx of internally displaced people that will increase the need and modify the buffer accordingly. [See Table 41.](#)
6. Finally, check the current stock levels for the balance. Subtract the stock balance from the estimated need.

Table 41. Example on How to Quantify Supplies Need

Estimating RUTF Needs	Estimating RUSF Needs
<ul style="list-style-type: none"> Estimate on average children 6-59 months will consume about 2 ½ sachets of RUTF a day. Each carton of RUTF contains 150 sachets of RUTF. <p>Example: Facility X has 20 children 6-59 months in OTP at the end of September 2018. The health facility will estimate the amount of RUTF needed for the next three months as follows:</p> <ul style="list-style-type: none"> RUTF required for 1 child with SAM for 1 months = 75 sachets (2.5 *30 days) RUTF required for 20 children for 3 months = 4500 sachets (75*20 children*3) Add 10% buffer stock = 4,950 (4500+450) Convert to cartons = 33 cartons (4,950/150) <p>Total amount to order for 3 months = 33 cartons of RUTF.</p> <p>Note: Always request full cartons of RUTF.</p>	<p>Children 6-59 months will consume 1 sachet of RUSF each month</p> <ul style="list-style-type: none"> Each carton of RUSF contains 150 sachets of RUSF. <p>Example: Facility X had 100 beneficiaries receiving TSFP services at the end of June 2018. The health facility will estimate the amount of RUSF needed for the next three months as follows:</p> <ul style="list-style-type: none"> RUSF required for 1 child with MAM for 1 month = 30 sachets RUSF required for 100 children for three months = 9,000 sachets (30*100 children*3) Add 10% buffer stock = 9,900 sachets (9000+900) Convert to cartons = 66 cartons (9,900/150) <p>Total amount to order for 3 months = 66 cartons of RUSF.</p> <p>Note: Always request full cartons of RUSF.</p>
	Estimating FBF Needs
	<ul style="list-style-type: none"> Children 6-59 months will consume 6 kg of FBF each month. <p>Example: Facility X had 100 children receiving TSFP services at the end of August 2018. The health facility will estimate the amount of FBF needed for the next three months as follows:</p> <ul style="list-style-type: none"> FBF required for 1 child with MAM for 1 month = 6 kg FBF required for 100 children for three months = 1,800 kg (6*100 beneficiaries*3) Add 10% buffer stock = 1,980 kg (1800+180) Convert kg into metric tonnes = 1.98 MT (1980/1000) <p>Total amount to order for 3 months = 1.98 MT of FBF</p> <ul style="list-style-type: none"> Pregnant or lactating women will consume 7.5 kg of FBF each month. <p>Facility X had 60 PLW receiving TSFP services at the end of August 2018. The health facility will estimate the amount of FBF needed for the next three months as follows:</p> <ul style="list-style-type: none"> FBF required for 1 pregnant or lactating woman in 1 month = 7.5 kg FBF required for 60 PLW for three months = 1,350 kg (7.5*60beneficiaries*3) Add 10% buffer stock = 1,485 kg (1350+135) Convert kg into metric tonnes = 1.485 MT (1980/1000) <p>Total amount to order for 3 months = 1.485 MT of FBF</p>

NOTE: in addition to specialised nutritious foods equipment and materials such as treatment cards, Multi-charts, and registration books should always be quantified and included in the request for supplies.

7.3.2 Place a Request

Supplies should be requested at the end of the month, in emergency situations, a request may be placed before the end of the month. To place a request for supplies, complete the “request for supplies” columns in the Monthly Supplies Report for SAM and MAM. [See Annex 27](#). Be sure to take note of the following:

1. Each item should be requested on its own row with a clear description of quantity needed, unit, and period of request in months.
2. Request for complete packs or cartons. For example, if you need 1000 sachets of RUTF, 1 carton contains 150 sachets, $1000/150 = 6.67$ cartons, therefore, request 7 cartons.

7.4 Receiving Supplies

At least two people should receive and check the new deliveries before they are put in the storage facility. The following should be checked:

- ◆ Delivery note, packing list, and contents against a copy of the requested items or products.
- ◆ Contents and quantity against the product list. Physically count to confirm.
- ◆ Outer and inner packaging for any signs of damage.
- ◆ Labels are legible and include complete information including the expiry date.
- ◆ Check the shelf life and expiry date. Do not accept products if the expiry date has passed.
- ◆ For equipment, check that all spare parts, accessories, instruction manuals, and warranty documents are included.
- ◆ Note any inconsistencies and report immediately to the next level.
- ◆ If the correct amount and condition of supplies matches the delivery note, receive and sign. Ensure that someone else verifies and signs a copy for record.
- ◆ File delivery documents together with the request documents.
- ◆ Store items and products immediately. Be sure to update the Bin Card and Stock Record Card with the quantities received. [See Annex 15 and 16](#) respectively.

7.5 Stock Management

7.5.1 Organising the Store

- ◆ SAM and MAM supplies should be stored in the general health facility store or pharmacy. Ensure that therapeutic and supplementary food products are not stored together with harmful chemicals.
- ◆ The storage facility should meet the following conditions:
 - ◆ The structure including the roof, ceiling, walls, and floor should be well maintained and lit.
 - ◆ Secure and always locked
 - ◆ Clean, tidy, and free of pests and rodents
 - ◆ Dry and at room temperature of 25-30°C
 - ◆ Well ventilated with adequate space for storage

-
- ◆ Has shelves or wooden pallets at least 10 cm from the wall
 - ◆ Not exposed to direct sunlight
 - ◆ Items and products should be arranged by expiry date to make it easy to follow the “First-Expiry, First-Out” (FEFO) principle.
 - ◆ Expired, damaged, or items no longer in use should be separated while awaiting disposal.
 - ◆ Expired items or products should be reported to the Food and Drug Authority through the Pharmaceutical Fund and Supply Agency for disposal according to rules and regulations.

7.5.2 Monitoring Stock

- ◆ The Bin Card and Stock Record Card should always be up-to-date with items or products received or issued.
- ◆ The HEW or store keeper should do a physical stock count of all items and products at the end of every month, before writing the monthly report.
- ◆ Be alert of stock outs, as soon noted, a request should be placed immediately.
- ◆ In case of excess stock, check the expiry date and report if the items or products need to be re-distributed to another location.
- ◆ Carry out an inventory of stock at least once a year.

7.5.3 Issuing Supplies

- ◆ Always record products or items issued from the store in the Bin Card and Stock Record Card.
- ◆ Record therapeutic or supplementary food commodities issued to beneficiaries in the treatment card and ration card.

7.6 Reporting Supplies

Report the essential SAM and MAM supplies at the end of every month using the Monthly Supplies Report for SAM and MAM. To prepare the report:

1. Confirm that each item or product is sorted by expiry date.
2. Carry-out a physical stock count.
3. Verify the following data from the Bin Card and Stock Record Card; received, issued, and losses or adjustments.

Note the following:

- ◆ The stock balance at the end of the previous month should always equal the stock balance at the beginning of the current month.
- ◆ The reason for losses or adjustment should always be noted in the remark’s column of the Monthly Supplies Report for SAM and MAM.

8 Monitoring, Reporting and Evaluation

This chapter describes the process and tools used to monitor and report SAM and MAM services. It also highlights on the methods used to evaluate access and coverage of SAM and MAM services.

8.1 Monitoring Community Outreach

Monitoring of community outreach activities is crucial in ensuring maximum coverage and access to SAM and MAM services. The following should be monitored at the community level:

- ◆ Training and capacity building activities conducted with the HDA/HDG.
- ◆ Frequency of community meetings and mobilisation activities, including the number and type of community groups and key stakeholders involved.
- ◆ Number of people screened and referred for treatment for acute malnutrition. Screening data should be compared with those who actually arrive to the health facility to obtain SAM and MAM services.
- ◆ Reasons for default and barriers to accessing care and treatment for SAM and MAM.

8.2 Monitoring Individual Patient Treatment

8.2.1 Registration Number

A registration number should be issued to each patient admitted to the OTP, SC, or TSFP. The registration number has the following information:

- ◆ The patient number which is assigned sequentially
- ◆ The service code indicating where treatment was initiated (i.e., OTP, SC or TSFP)
- ◆ The patient's medical record number.
- ◆ The same registration number should be maintained even if the patient is transferred or referred for services in another health facility.
- ◆ The registration number should be used in all monitoring and referral documents pertaining to the patient. These include the Multi-chart, treatment and follow-up cards, ration card, registration books, and referral slip.

8.2.2 Individual Treatment and Follow-up Card

Outpatient Therapeutic Programme (OTP) Treatment and Follow-up Card

The OTP Treatment and Follow-up Card should be issued to each patient admitted to the OTP. [See Annex 17](#). The card is used to record information on the initial medical history, physical examination, routine medications, and follow-up visits.

Stabilisation Centre (SC) Multi-chart

The SC Multi-chart should be issued to each patient admitted to the SC. [See Annex 18](#). The Multi-chart is used to record information on the initial care, medical and physical assessment, daily surveillance, and treatment outcomes.

Targeted Supplementary Feeding Programme (TSFP) Treatment and Follow-up Card

Patients receiving treatment for MAM should be issued a TSFP Treatment and Follow-up Card. The card is used to record anthropometric information, rations, and routine medications provided to the patient. [See Annex 19](#) and [Annex 20](#) for TSFP cards for children and PLW respectively.

TSFP Identification Card

A TSFP Identification Card should be issued to every patient admitted and receiving MAM treatment. The patient keeps the card and uses it for identification during follow-up visits at the health facility. [See Annex 21](#) for the TSFP Identification Card for Admitted Patients.

Filing System

An efficient system of filling the SC Multi-charts, OTP and TSFP cards should be maintained during admission, feeding, drug administration and follow-up of children. Each health facility should have two files for each service (i.e., SC, OTP or TSFP). [Table 42](#) below summarizes how the Multi-charts and treatment and follow-up cards should be filed.

Table 42. The Filing System used in SC, OTP and TSFP

File 1 Currently in Treatment	File 2 Discharged
<ul style="list-style-type: none"> ■ Currently in SC, OTP or TSFP ■ Absentees ■ Transfers from one TSFP to another/or from one OTP to another 	<ul style="list-style-type: none"> ■ Cured * ■ Defaulters ** ■ Non-responder ■ Deaths ■ Others (i.e., pregnant women who delivered or lactating women whose infant is ≥ 6 months)

* A separate file may be required for discharged cured cases due to the large number of cards.

** If defaulters return for treatment, monitoring continues with the same card/registration number.

8.2.3 Registration Books

The registration books provide a summary of SAM and MAM treatment. They help with the compilation of the Monthly Statistics Report for SAM and MAM. The registration books include information on the patient admission criteria and discharge outcomes. There are three separate registration books used for the management of SAM and MAM:

- ◆ Registration book for SAM Treatment. [See Annex 23](#).
- ◆ Registration book for MAM treatment – Children 6-59 Months and Older. [See Annex 24](#).
- ◆ Registration book for MAM Treatment – PLW. [See Annex 25](#).

8.2.4 Referral Slip

If a patient is referred from OTP to SC and vice versa, or from TSFP to OTP, a referral slip should be given to the caregiver with instructions on how and when to go where. The referral slip should also provide information on medication or treatment provided prior to referral. [See Annex 22](#).

8.3 Reporting

- ◆ The Monthly Statistics Report for SAM and MAM is used to record data on SAM and MAM admissions, transfers, and discharges. [See Annex 26](#).
- ◆ The Monthly Supplies Report for SAM and MAM captures data on supply management including request for supplies. [See Annex 27](#).
- ◆ Each health facility, woreda, zone and region should submit the monthly statistics report and monthly supplies report to the next level for consolidation and further analysis.

[Table 43](#) below provides a summary of the entry and exit categories in SAM and MAM treatment.

Table 43. Summary of Entry and Exit Categories

SC for the Management of SAM with Medical Complications	OTP for the Management of SAM without Medical Complications	TSFP for the Management of MAM
Entry Categories		
<p>New admission: New case who meets the admission criteria for SC.</p> <p>Readmission: Re-admitted for treatment in SC due to <u>relapse</u> or a <u>returned defaulter</u> who meets the admission criteria.</p> <p>Relapse: Cured within the past 3 months and now meets the admission criteria for SC.</p> <p>Returned defaulter: Defaulted within the past 3 weeks and has returned to continue treatment in SC. A returned defaulter should be re-admitted if they meet the admission criteria.</p> <p>Transfers in: Referred to the SC because condition deteriorated in OTP or has moved in from another facility where s/he was receiving care in SC.</p>	<p>New admission: New case who meets the admission criteria for OTP.</p> <p>Readmission: Re-admitted for treatment in OTP due to <u>relapse</u> or a <u>returned defaulter</u> who meets the admission criteria.</p> <p>Relapse: Cured within the past 3 months and now meets the admission criteria for OTP. Relapse cases require special attention therefore should be referred to start treatment in SC.</p> <p>Returned defaulter: Defaulted within the past 3 months and has returned to continue treatment in OTP. A returned defaulter should be re-admitted if they meet the admission criteria.</p> <p>Transfers in: Continuing treatment in OTP after stabilisation in SC or has moved in from another facility where s/he was receiving OTP.</p>	<p>New admission: New case who meets the admission criteria for TSFP.</p> <p>Readmission: Re-admitted for treatment in TSFP due to <u>relapse</u> or a <u>returned defaulter</u> who meets the admission criteria.</p> <p>Relapse: Cured within the past 3 months and now meets the admission criteria for TSFP.</p> <p>Returned defaulter: Defaulted within the past 3 months and has returned to continue treatment in TSFP. A returned defaulter should be re-admitted if they meet the admission criteria.</p> <p>Transfers in: Has moved in from another facility where s/he was receiving TSFP.</p>
Exit Categories		
<p>Cured: Has reached the discharge criteria for SAM treatment, i.e., the special cases that were not referred to OTP.</p> <p>Died: Dies while receiving treatment in the SC.</p> <p>Defaulted: Absent for two consecutive days. Default should be confirmed.</p> <p>Non-responder: Patient who remained in treatment in the SC does not reach the SAM discharge criteria after 16 weeks (4 months) in treatment.</p> <p>Stabilised: Condition has stabilised and referred to continue treatment in OTP.</p> <p>Transfer out: Moved to another facility for further medical care or moved out to receive care in another SC.</p>	<p>Cured: Has reached the discharge criteria for SAM treatment.</p> <p>Died: Dies while receiving treatment in the OTP.</p> <p>Defaulted: Absent for two consecutive visits. Default should be confirmed.</p> <p>Non-responder: Does not reach the SAM discharge criteria after 16 weeks (4 months) in treatment.</p> <p>Transfer out: Condition has deteriorated or not responding to treatment according to action protocol and referred for treatment in the SC or moved out to receive OTP in another facility.</p>	<p>Cured: Has reached the discharge criteria for MAM treatment.</p> <p>Died: Dies while receiving treatment in the TSFP.</p> <p>Defaulted: Absent for two consecutive visits. Default should be confirmed.</p> <p>Non-responder: Does not reach the MAM discharge criteria after 16 weeks (4 months) in treatment.</p> <p>Transfer out: Condition has deteriorated to SAM and referred for OTP or moved out to receive TSFP in another facility.</p> <p>Others: Pregnant woman who has delivered or a lactating woman whose infant is 6 months</p>

8.4 Monitoring Programme Performance

The performance of SAM and MAM services can be analysed from the monthly report and outcomes compared to international Sphere standards.

8.4.1 Programme Indicators

The outcomes are calculated as a proportion of SAM or MAM programme discharges over a period, usually monthly, quarterly, or annually. When calculating SAM outcomes, the SC and OTP discharges should be combined for a comprehensive analysis of the SAM programme performance.

The following programme outcomes are monitored.

1. **Cure rate:** Proportion who are discharged from SAM or MAM treatment having reached the cure discharge criteria.

$$\text{Cure rate (\%)} = \frac{\text{Total discharged cured}}{\text{Cured} + \text{Died} + \text{Defaulted} + \text{Non responders}} \times 100$$

2. **Default rate:** Proportion discharged having defaulted.

$$\text{Default rate (\%)} = \frac{\text{Total discharged defaulted}}{\text{Cured} + \text{Died} + \text{Defaulted} + \text{Non responders}} \times 100$$

3. **Death rate:** Proportion discharged having died while registered for SAM or MAM treatment.

$$\text{Death rate (\%)} = \frac{\text{Total discharged died}}{\text{Cured} + \text{Died} + \text{Defaulted} + \text{Non responders}} \times 100$$

4. **Non-responder:** Proportion discharged having not achieve the cure discharge criteria.

$$\text{Non-responder rate (\%)} = \frac{\text{Total discharged non responder}}{\text{Cured} + \text{Died} + \text{Defaulted} + \text{Non responders}} \times 100$$

8.4.2 Other Service Indicators

Average length of stay (LOS): The number of days that a patient spends in treatment from admission to discharge. LOS is only calculated for patients cured.

$$\text{Average Length of Stay (LOS)} = \frac{\text{Sum LOS}}{\text{Number of cards or cases in the sample}}$$

Average Weight Gain (AWG): The rate of weight gain per kilogram of body weight per day. AWG is only calculated for patients discharged cured.

$$\text{*Weight Gain} = \frac{\text{Discharge weight in grams} - \text{Minimum weight in grams}}{\text{Minimum weight in kg} \times \text{number of days between minimum weight and discharge day}}$$

$$\text{Average Weight Gain (AWG)} = \frac{\text{Sum of weight gains (in grams per kg bodyweight per day)}}{\text{Number of cards or cases in the sample}}$$

**Note: When calculating the LOS and weight gain, separate the OTP cards in two group: those who had bilateral pitting oedema, and those with wasting (MUAC + WFH).*

8.4.3 Outcome Indicators

The table below provides the international reference standards for MAM and SAM outcome indicators.

Table 44. Outcome Indicators


Indicator	Management of SAM	Management of MAM
Cured	> 75%	> 75%
Defaulted	< 15%	< 15%
Died	< 10%	< 3%
Non-responder	Not stated	Not stated
Length of stay	Approximately 60 days	Approximately 90 days
Average Weight Gain	*>5 g/kg/day	Not applicable

*A weight gain of 5g/kg/day is a suggested guideline (WHO 2005) for community-based programmes. Weight gains may vary for patients with wasting and bilateral pitting oedema.

8.4.4 Data Compilation and Quality Assurance

- ◆ Following the compilation and analysis of the Monthly Statistics Report for Acute Malnutrition, the reports should be sent to the Woreda health office for entry into the Health Management Information System (HMIS), and then to Zonal, Regional, and Federal levels. [Figure 9](#) illustrates the data flow.
- ◆ The compilation sheets should be used for compiling data from various sites. Excel or other spreadsheet software can be used to facilitate calculations, minimise errors, and speed-up data processing.

Figure 9. Acute Malnutrition Data Flow



Facility service providers, managers, and supervisors should consider the following aspects to ensure data quality. (See Table 45).

Table 45. Data Quality Assurance

Accuracy	Whether the data recorded and reported are correct
Reliability	Whether the data are consistent. This can be checked by comparing the data from previous months or the same month in previous years.
Completeness	Whether all necessary data elements are filled. At the cluster health centre level or above, the completeness of health facility reports (i.e., facility reporting rate) give an indication of programme functionality and performance.
Timeliness	Whether all expected reports are ready and submitted within a specified time frame.
Legibility	Whether the data written, transcribed and/or printed is readable.
Accessibility	Whether the needed data is available for individual SAM or MAM patient care, programme planning, or decision making.

8.5 Supportive Supervision

The Woreda health offices, ZHD and RHBs should conduct regular supportive supervision visits to health facilities. The supervisor(s) should use the supervisor scorecard to monitor the quality of service delivery. **See Annex 28.** The following should be reviewed during the visit:

- ◆ Adherence to admission and discharge criteria for SAM and MAM
- ◆ Accurate patient tracking, completion of treatment cards, registers, and report forms
- ◆ Proper completion of the treatment and monitoring cards
- ◆ Adherence to the medical treatment and nutrition rehabilitation protocols
- ◆ Progress of individual children, checking for consistent weight gain
- ◆ Routine and active follow-up and referral of cases
- ◆ SAM and MAM supply management
- ◆ Linkage and referral with other health and nutrition interventions

8.6 Coverage Assessment

Coverage assessments is specific to the evaluation of the acute malnutrition service delivery. It identifies the proportion of cases with acute malnutrition who are receiving services within a given area (Kebele, Woreda, Zone or Region).

For example:

$$\text{Treatment Coverage for Acute Malnutrition} = \frac{\text{Number receiving treatment for SAM+MAM}}{\text{Total number of SAM+MAM cases in the area}}$$

Coverage assessments can be conducted using the following techniques and methods:

- ◆ The Simplified Lot Quality Assurance Sampling Evaluation of Access and Coverage (SLEAC) is a low-cost classification-based version of Centric Systematic Area Sampling (CSAS). SLEAC is used together with the Semi Quantitative Evaluation of Access and Coverage (SQUEAC) which uses a semi-quantitative approach to investigate factors that influence the coverage of services. More information on SQUEAC and SLEAC is available at: <https://www.fantaproject.org/monitoring-and-evaluation/squeac-sleac>.
- ◆ The Simple Spatial Survey Method (S3M) is an improved version of the CSAS. The advantage of S3M is that it can be used for wide scale assessment of coverage at a relatively low cost. The S3M is also easy to understand and apply. More information on S3M are available at: http://www.validinternational.org/coverage/workshop/articles_files/pictureBookS3M.pdf



References

1. Briend, A.; Mwangome, M.K.; and Berkley, J.A. 2017. "[Using Mid-Upper Arm Circumference to Detect High-Risk Malnourished Patients in Need of Treatment.](#)" Preedy, V.R. and Patel, V.B. (eds.). Handbook of Famine, Starvation, and Nutrient Deprivation. DOI 10.1007/978-3-319-40007-5_11-1.
2. Dessie, M. et al. 2015. Community Engagement for CMAM. Coverage Monitoring Network. Available at: <http://www.coverage-monitoring.org/wp-content/uploads/2015/07/Technical-brief-Community-Engagement-for-CMAM-20151.pdf>.
3. Federal Ministry of Health (FMOH). 2015. Health Sector Transformation Plan (HSTP) 2015/16 – 2019/20. Addis Ababa: FMOH. Available at: <http://www.moh.gov.et/hu/web/guest/-/health-sector-transformation-plan?inheritRedirect=true>
4. Food and Nutrition Technical Assistance III Project (FANTA). 2018. Training Guide for Community-Based Management of Acute Malnutrition (CMAM). Washington DC: FHI 360/FANTA. Available at: <https://www.fantaproject.org/focus-areas/nutrition-emergencies-mam/cmam-training>
5. Food, Medicine and Health Administration and Control Authority of Ethiopia (FMHACA). 2014. Standard Treatment Guidelines for General Hospital. Addis Ababa: FMHACA.
6. FMHACA. 2014. *Standard Treatment Guidelines for Primary Hospital*. Addis Ababa: FMHACA.
7. FMHACA. 2014. *Standard Treatment Guidelines for Primary Health Centres*. Addis Ababa: FMHACA.
8. FMOH. 2016. *Health Sector Transformation Plan 2015/16-2019/20*. Addis Ababa: FMOH.
9. FMOH. 2010. *Integrated Management of Newborn and Childhood Illness (IMNCI)*. Addis Ababa: FMOH.
10. FMOH. 2012. *Management of Moderate Acute Malnutrition Guidelines*. Addis Ababa: FMOH.
11. FMOH. 2018. *National Guidelines for Comprehensive HIV Prevention, Care and Treatment*. Addis Ababa: FMOH.
12. FMOH. 2017. *National Guidelines for Management of Tuberculosis (TB), Drug-Resistant TB and Leprosy in Ethiopia*. Addis Ababa: FMOH.
13. FMOH. 2016. *National Guidelines on Adolescent, Maternal Infant and Young Child Nutrition*. Addis Ababa: FMOH.
14. FMOH. 2017. *National Implementation Guidelines for Integrated Community Case Management of Childhood Illnesses and Newborn Care*. Addis Ababa: FMOH.
15. FMOH. 2016. *National Nutrition Programme (NNP) II 2016-2020*. Addis Ababa: FMOH.
16. FMOH. 2007. *National Protocol for the Management of Severe Acute Malnutrition*. Addis Ababa: FMOH.

-
17. Global Nutrition Cluster. 2015. *CMAM Caseload Calculations (SAM and MAM Treatment)*. Available at: <http://nutritioncluster.net/resources/caseload-targets-supplies-calculator/>
 18. Global Nutrition Cluster. 2017. *Moderate Acute Malnutrition: A Decision Tool for Emergencies*. Available at: <http://nutritioncluster.net/?get=002086%7C2014/07/MAM-Decision-Tool-final-June-2014-corrected.pdf>
 19. Kerac, M. et al. 2014. "The interaction of malnutrition and neurologic disability in Africa." *Seminars in Pediatric Neurology*, 21 (1). pp. 42-9. ISSN 1071-9091 DOI. Available at: <http://researchonline.lshtm.ac.uk/1924646/>
 20. Sphere Association. 2018. *The Sphere Handbook: Humanitarian Charter and Minimum Standards in Humanitarian Response, Fourth Edition*. Geneva: Sphere. Available at: <http://www.spherehandbook.org/en/contents-3/>
 21. World Health Organisation (WHO). 2013. *Guideline: Nutrition Care and Support for Patients with Tuberculosis*. Geneva: WHO. Available at: http://apps.who.int/iris/bitstream/handle/10665/94836/9789241506410_eng.pdf;jsessionid=7AF68F46055D29DFC3CABBFD623638B2?sequence=1
 22. WHO. 2013. *Guideline: Updates on the Management of Severe Acute Malnutrition in Infants and Children*. Geneva: WHO. Available at: http://www.who.int/nutrition/publications/guidelines/updates_management_SAM_infantandchildren/en
 23. WHO. 2003. *Guidelines for the Inpatient Treatment of Severely Malnourished Children*. Geneva: WHO. Available at: <http://www.who.int/nutrition/publications/severemalnutrition/9241546093/en/>
 24. WHO. 1999. *Management of Severe Malnutrition: A Manual for Physicians and Other Senior Health Workers*. Geneva: WHO. Available at: www.who.int/nut/publications
 25. WHO. 2013. *Pocket Book of Hospital Care for Children: Guidelines for Management of Common Childhood Illnesses. 2nd Edition*. Available at: http://www.who.int/maternal_child_adolescent/documents/child_hospital_care/en/
 26. WHO and UNICEF. 2009. *WHO Child Growth Standards and Identification of Infants and Children with Severe Acute Malnutrition: A Joint Statement*. Available at: <http://www.who.int/nutrition/publications/severemalnutrition/9789241598163/en/>



Annex 1. Home Visit Checklist

Health Extension Worker (HEW)/ Community Worker's Name: _____

Name of Patient: _____

Date of Visit: _____

Kebele: _____

Gott: _____

Feeding Practices

Is the specialised nutritious food (i.e., RUTF, RUSF or FBF) present in the home?	Yes	No
If not, where is the specialised nutritious food?		
Is the available specialised nutritious food enough to last until the next OTP or TSFP follow-up visit?	Yes	No
Is the specialised nutritious food being shared or eaten only by the patient?	Shared	Patient only
Is the patient given other foods in addition to the specialised nutritious food?	Yes	No
If yes, what type of food?		
How many times in a day does the patient eat?		
Does someone help/encourage the patient to eat?	Yes	No
What does the caregiver do if the patient does not want to eat?		
If 0–23 months old, is the child currently breastfeeding?	Yes	No

Caring Practices (Children 0–59 months)

Are both parents alive and healthy?	Yes	No
Who cares for the child during the day?		
Does the child look washed and clean?	Yes	No

Health and WASH

Is clean water available to the household?	Yes	No
What is the household's main source of water?		
Is soap available for use by household members?	Yes	No
Does the patient wash hands and face before feeding?	Yes	No
Is food covered and protected from flies?	Yes	No
What action does the caregiver take when the child has diarrhoea?		

Food Security

Does the household currently have food available?	Yes	No
What is the main household source of income?		
Is the household registered for PSNP?	Yes	No
Does the household receive any other form of external support?	Yes, specify: _____	No



Annex 2. Management of Acute Watery Diarrhoea (AWD) and SAM

Key Messages:

- ◆ All patients with SAM and suspected AWD must be treated at a Cholera Treatment Centre (CTC) as rehydration should be addressed before nutrition care and treatment is initiated.
- ◆ Nutritional status of patients with AWD should be assessed as management differs if the patient has SAM.
- ◆ Patients with SAM have different physiology so they must be rehydrated slowly. IV fluids should only be used for SAM patients in shock because of the high risk of fluid overload and heart failure.
- ◆ Children with SAM and AWD must be treated for dehydration using low-osmolarity Oral Rehydration Solution (ORS). Do not use ReSoMal.
- ◆ During rehydration, closely monitor signs of fluid overload.
- ◆ Re-assess and re-classify nutritional status after rehydration. Adjust management if needed.
- ◆ Patients with SAM and AWD should be treated with the same therapeutic feeds, following the feeding protocol for patients with SAM and medical complications.
- ◆ As soon as the patient has recovered from AWD, nutritional status should be reassessed and the child referred to the Stabilisation Centre (SC).
- ◆ Breastfed infants should continue with breastfeeding as it the safest source of nutrition.

Principles of Management

Diagnosis

- ◆ A diagnosis of AWD is considered if:
 - ◆ There is evidence of high output diarrhoea (one stool an hour).
 - ◆ Diarrhoea appears pale and straw-coloured.
 - ◆ Diarrhoea may be accompanied by vomiting and nausea.
 - ◆ A patient's family members have been diagnosed with or have been suspected of having AWD.
- ◆ A diagnosis of SAM should be made before starting treatment of AWD.

In children 6-59 months check:

- ◆ Is there bilateral pitting oedema?
- ◆ Is MUAC < 11.5cm?
- ◆ Is WFH < -3 z-score?

.....

In infants 0-6 months check:

- ◆ Is there bilateral pitting oedema?
- ◆ Is WFL < -3 z-score?
- ◆ Does the infant have recent weight loss or failure to gain weight?
- ◆ For a patient with SAM and diarrhoea, do a rapid diagnostic test (Crystal VC Dipstick), a screening test for AWD, if available. Collect stool samples for confirmatory diagnostic testing.

Note: Diagnosis of SAM in patients with AWD can be difficult as dehydration can cause loss of body weight and affect the MUAC and WFH/WFL measurement. Weight and MUAC must be reassessed after rehydration and at discharge from the CTC to confirm the nutritional status and refer for appropriate nutrition care and treatment.

- ◆ Diagnosis of dehydration in a patient with SAM and AWD should be considered if:
 - ◆ There is a clear history of a recent change in the patient's appearance.
 - ◆ The eyes are sunken since the start of diarrhoea or vomiting (history confirmed by mother or caretaker).
- ◆ Diagnosis of shock in a patient with SAM should be considered if the patient shows signs of weak or absent radial or femoral pulse, cool or cold hands and feet, low or unreportable blood pressure, or if there is loss of consciousness.

Management


- ◆ AWD results in extensive loss of fluid and electrolytes and can kill rapidly so rehydration should be addressed before nutrition care and treatment initiated. Do not keep patients with AWD in OTP or SC, immediately refer them to a CTC for rehydration and to avoid cross infection.
- ◆ Management depends on the dehydration status of the patient. See steps 1 to 7 below for details.
- ◆ Oral rehydration with low osmolarity ORS should be used. ReSoMal should not be used to rehydrate patients with SAM and AWD.
- ◆ Patients with SAM and AWD, who present with dehydration but are not in shock, should be rehydrated slowly using ORS.
- ◆ Because of the high risk of fluid overload and heart failure, IV hydration should be avoided in children with SAM and AWD unless if the patient is in shock.
- ◆ Therapeutic foods already contain adequate zinc, therefore children with SAM and AWD receiving F-75, F-100 or RUTF should not receive additional zinc supplements.
- ◆ Breastfed children should continue to breastfeed.

STEPS: Management of AWD with SAM

STEP 1: Determine nutritional status

- ◆ Does the patient have SAM?
 - ◇ If no, use the Standard Rehydration Procedures for AWD and refer to the AWD Clinical Guidelines
 - ◇ If yes, follow step 2 (assess for dehydration and shock in the SAM patient)

STEP 2: Assess dehydration and shock in patients with SAM



STEP 3: Treatment for non-dehydrated patients with SAM and AWD

- ◆ Continue breastfeeding and age-appropriate food.
- ◆ Administer ORS to replace ongoing losses:
 - ◇ If < 2 years old, 50-100 ml per loose stool
 - ◇ If > 2 years old, 100-200 ml per loose stool
- ◆ As soon as the patient is stable, refer to the SC for the management of SAM.

STEP 4: Treatment for dehydrated patients with SAM and AWD - CONSCIOUS and able to drink.

- ◆ 5 ml/kg of ORS every 30 minutes for the first 2 hours
- ◆ 5-10 ml/kg per hour of ORS, alternating with F-75, for a maximum of 10 hours or until the fluid deficit is corrected

-
- ◆ Adjust ORS intake during rehydration phase to compensate for on-going fluid loss in high-output stooling
 - ◆ After rehydration give:
 - ◇ If < 2 years and wasted, 50-100 ml of ORS orally after each watery stool.
 - ◇ If > 2 years and wasted, 100-200 ml of ORS orally after each loose stool.
 - ◇ If the child has bilateral pitting oedema, give 30 ml of ORS orally per each loose stool.
 - ◆ If the patient cannot drink adequately or is unable to drink, administer ORS via NGT.

STEP 5: Treatment for dehydrated patients with SAM and AWD- in shock

- ◆ If the patient has the following signs of shock:
 - ◇ Unresponsive or semi-conscious
 - ◇ Absent or weak pulse
 - ◇ Cold hands and feet
 - ◇ Continuing vomiting after introduction of NGT OR
 - ◇ Presence of abdominal distension
- ◆ Give IV treatment as follows:
 - ◇ Give IV Ringer lactate with 5% dextrose.
 - ◇ 15 ml/kg/h for the first hour then reassess, if there is improvement (a decrease in respiratory and pulse rates), same amount repeated for another one hour.
 - ◇ After 2 hours of IV fluids, give 10 ml/kg per hour of ORS orally or via NGT until the deficit is corrected or until the patient is fully rehydrated.
 - ◇ Adjust IV flow rate during the rehydration phase to compensate for ongoing fluid loss in
 - ◇ high-output stooling.
- ◆ Check every 10 minutes for heavy or laboured breathing.
- ◆ In case of presence of one or more signs of fluid overload or cardiac failure (heavy, laboured breathing, engorged jugular vein pressure or increased oedema) then stop the IV and consult a physician.

STEP 6: Ongoing monitoring

Continue to check the patient's status. Reassess the following after 1 hour:

- ◆ If the breathing status worsens, then stop IV infusion.
- ◆ If you do not see improvement, consider septic shock.
- ◆ If you do see improvement, continue the same amount of IV fluid for the next one hour.

STEP 7: Rehydration complete

- ◆ Rehydration is complete when:
 - ◆ Patient is no longer thirsty.
 - ◆ Urine production has normalised.
 - ◆ Other signs of dehydration have resolved.
- ◆ Once hydration is re-established, measure MUAC. If MUAC is still less than 11.5 cm, continue with treatment for non-dehydrated children with SAM.
- ◆ Initiate feeding and treat medical complications such as hypothermia and/or hypoglycaemia.



Annex 3. Antibiotics for Patients with SAM

IF:	GIVE:		
NO MEDICAL COMPLICATIONS	Amoxicillin oral: 25 mg/kg every 12 hours for 5 days or until referral to OTP		
MEDICAL COMPLICATIONS (shock, hypoglycaemia, hypothermia, dermatosis with raw skin/fissures, respiratory or urinary tract infections, or lethargic/sickly appearance)	* Gentamicin IV or IM (5 mg/kg) once daily for 7 days, plus: <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days</td> <td>Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days</td> </tr> </table>	Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days	Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days
Ampicillin IV or IM (50 mg/kg) every 6 hours for 2 days	Followed by: **Amoxicillin oral (25 mg/kg), every 12 hours for 5 days		
Resistance to Amoxicillin and Ampicillin and Presence of Medical Complications	In the case of sepsis or septic shock , give: IV/IM ceftriaxone (children or infants over 1 month of age (50 mg/kg every 8 to 12 hours) + oral ciprofloxacin (5 to 15 mg/kg 2 times per day) If suspected staphylococcal infection , add: IV/IM cloxacillin (12.5–50 mg/kg/dose four times a day, depending on the severity of the infection).		
Specific Infection Requires an Additional Antibiotic	Add specific antibiotic as per the Ethiopia Standard Treatment Guidelines.		
If HIV-positive or Child who is HIV Exposed.	Cotrimoxazole oral according to the National Guidelines for Comprehensive HIV Prevention, Treatment and Care.		

*If the child is not passing urine, gentamicin may accumulate in the body and cause deafness. Do not give the second dose until the child is passing urine

**If amoxicillin is not available, give ampicillin, 50 mg/kg orally every 6 hours for 5 days

Specific Formulations and Body Weight Ranges for Antibiotics for SAM Children in SC

ANTIBIOTIC	ROUTE/DOSE/ FREQUENCY/ DURATION	FORMULATION
Amoxicillin	Oral: 25 mg/kg body weight every 12 hours for 5 days	Syrup, 125 mg/5 ml
		Syrup, 250 mg/5 ml
Ampicillin	IV or IM: 50 mg/kg every 6 hours for 2 days	Vial of 500 mg mixed with 2.1 ml sterile water to give 500 mg /2.5 ml
Metronidazole	Oral: 5mg/kg twice a day for 4 days maximum	Suspension, 40mg/ml
Benzylpenicillin	IV or IM: 50,000 units/kg body weight every 6 hours for 5 days	IV: vial of 600 mg mixed with 9.6 ml sterile water to give 1,000,000 IU/10 ml
		IM: vial of 600 mg mixed with 1.6 ml sterile water to give 1,000,000 IU/2 ml

ANTIBIOTIC	ROUTE/DOSE FREQUENCY/ DURATION	FORMULATION	DOSES FOR SPECIFIC BODY WEIGHTS (Use closest weight)									
			3 kg	4 kg	5 kg	6 kg	7 kg	8 kg	9 kg	10 kg	11 kg	12 kg
Gentamicin	IV or IM: 5 mg/kg once daily for 7 days	IV/IM: vial containing 20 mg (2 ml at 10 mg/ml), undiluted	2.25 ml	3.0 ml	3.75 ml	4.5 ml	5.25 ml	6.0 ml	6.75 ml	7.5 ml	8.25 ml	9.0 ml
		IV/IM: vial containing 80 mg (2 ml at 40 mg/ml) mixed with 6 ml sterile water to give 80 mg/8 ml	2.25 ml	3.0 ml	3.75 ml	4.5 ml	5.25 ml	6.0 ml	6.75 ml	7.5 ml	8.25 ml	9.0 ml
		IV/IM: vial containing 80 mg (2 ml at 40 mg/ml), undiluted	0.5 ml	0.75 ml	0.9 ml	1.1 ml	1.3 ml	1.5 ml	1.7 ml	1.9 ml	2.0 ml	2.25 ml

Specific Formulations and Body Weight Ranges for Antibiotics for SAM Children in SC

Weight of Child	Dose of Iron Syrup: Ferrous Fumarate 100mg per 5ml (20mg elemental iron per ml)
3 up to 6 kg	0.5 ml
6 up to 10 kg	0.75 ml
10 kg to 15 kg	1 ml



Annex 4. F-75 Reference Tables for Stabilisation Phase (Severe Wasting)

Weight of Child (kg)	Volume of F-75 per Feed (ml) ^a			Daily Total (130 ml/kg)	80% of Daily Total ^a (Minimum)
	Every 2 Hours ^b (12 Feeds)	Every 3 Hours ^c (8 Feeds)	Every 4 Hours (6 Feeds)		
2.0	20	35	45	260	210
2.2	25	35	50	286	230
2.4	25	40	50	312	250
2.6	30	40	55	338	270
2.8	30	45	60	364	290
3.0	35	50	65	390	310
3.2	35	50	70	416	335
3.4	35	55	75	442	355
3.6	40	60	80	468	375
3.8	40	60	80	494	395
4.0	45	65	85	520	415
4.2	45	70	90	546	435
4.4	50	70	95	572	460
4.6	50	75	100	598	480
4.8	50	80	105	624	500
5.0	55	80	110	650	520
5.2	55	85	115	676	540
5.4	60	90	115	702	560
5.6	60	90	120	728	580
5.8	65	95	125	754	605
6.0	65	100	130	780	625
6.2	65	100	135	806	645
6.4	70	105	140	832	665
6.6	70	105	145	858	685
6.8	75	110	145	884	705
7.0	75	115	150	910	730
7.2	80	115	155	936	750
7.4	80	120	160	962	770
7.6	80	125	165	988	790
7.8	85	125	170	1014	810
8.0	85	130	175	1040	830
8.2	90	135	180	1066	855

8.4	90	135	180	1092	875
8.6	95	140	185	1118	895
8.8	95	145	190	1144	915
9.0	100	145	195	1170	935
9.2	100	150	200	1196	955
9.4	100	155	205	1222	980
9.6	105	155	210	1248	1000
9.8	105	160	210	1274	1020
10.0	110	165	215	1300	1040
10.2	110	165	220	1326	1060
10.4	115	170	225	1352	1080
10.6	115	170	230	1378	1100
10.8	115	175	235	1404	1125
11.0	120	180	240	1430	1145
11.2	120	180	245	1456	1165
11.4	125	185	245	1482	1185
11.6	125	190	250	1508	1205
11.8	130	190	255	1534	1225
12.0	130	195	260	1560	1250
12.2	130	200	265	1586	1270
12.4	135	200	270	1612	1290
12.6	135	205	275	1638	1310
12.8	140	210	275	1664	1330
13.0	140	210	280	1690	1350
13.2	145	215	285	1716	1375
13.4	145	220	290	1742	1395
13.6	145	220	295	1768	1415
13.8	150	225	300	1794	1435
14.0	150	230	305	1820	1455
14.2	155	230	310	1846	1475
14.4	155	235	310	1872	1500
14.6	160	235	315	1898	1520
14.8	160	240	320	1924	1540
15.0	165	245	325	1950	1560
15.2	165	245	330	1976	1580
15.4	165	250	335	2002	1600
15.6	170	255	340	2028	1620
15.8	170	255	340	2054	1645
16.0	175	260	345	2080	1665
16.2	175	265	350	2106	1685
16.4	180	265	355	2132	1705

16.6	180	270	360	2158	1725
16.8	180	275	365	2184	1745
17.0	185	275	370	2210	1770
17.2	185	280	375	2236	1790
17.4	190	285	375	2262	1810
17.6	190	285	380	2288	1830
17.8	195	290	385	2314	1850
18.0	195	295	390	2340	1870
18.2	195	295	395	2366	1895
18.4	200	300	400	2392	1915
18.6	200	300	405	2418	1935
18.8	205	305	405	2444	1955
19.0	205	310	410	2470	1975
19.2	210	310	415	2496	1995
19.4	210	315	420	2522	2020
19.6	210	320	425	2548	2040
19.8	215	320	430	2574	2060
20.0	215	325	435	2600	2080

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Give 2-hourly feeds for at least the first day. When there is little or no vomiting, moderate diarrhoea (< 5 watery stools per day), and the child finishes most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds, if there is no vomiting, less diarrhoea, and the child finishes most feeds, change to 4-hourly feeds.

Annex 5. F-75 Reference Tables for Stabilisation Phase (Bilateral Pitting Oedema +++)

Weight with +++ Oedema (kg)	Volume of F-75 per Feed (ml) ^a			Daily Total (100 ml/kg)	80% of Daily Total (Minimum)
	Every 2 Hours ^b (12 Feeds)	Every 3 Hours ^c (8 Feeds)	Every 4 Hours (6 Feeds)		
3.0	25	40	50	300	240
3.2	25	40	55	320	255
3.4	30	45	55	340	270
3.6	30	45	60	360	290
3.8	30	50	65	380	305
4.0	35	50	65	400	320
4.2	35	55	70	420	335
4.4	35	55	75	440	350
4.6	40	60	75	460	370
4.8	40	60	80	480	385
5.0	40	65	85	500	400
5.2	45	65	85	520	415
5.4	45	70	90	540	430
5.6	45	70	95	560	450
5.8	50	75	95	580	465
6.0	50	75	100	600	480
6.2	50	80	105	620	495
6.4	55	80	105	640	510
6.6	55	85	110	660	530
6.8	55	85	115	680	545
7.0	60	90	115	700	560
7.2	60	90	120	720	575
7.4	60	95	125	740	590
7.6	65	95	125	760	610
7.8	65	100	130	780	625
8.0	65	100	135	800	640
8.2	70	105	135	820	655
8.4	70	105	140	840	670
8.6	70	110	145	860	690
8.8	75	110	145	880	705
9.0	75	115	150	900	720
9.2	75	115	155	920	735

9.4	80	120	155	940	750
9.6	80	120	160	960	770
9.8	80	125	165	980	785
10.0	85	125	165	1000	800
10.2	85	130	170	1020	815
10.4	85	130	175	1040	830
10.6	90	135	175	1060	850
10.8	90	135	180	1080	865
11.0	90	140	185	1100	880
11.2	95	140	185	1120	895
11.4	95	145	190	1140	910
11.6	95	145	195	1160	930
11.8	100	150	195	1180	945
12.0	100	150	200	1200	960
12.2	100	155	205	1220	975
12.4	105	155	205	1240	990
12.6	105	160	210	1260	1010
12.8	105	160	215	1280	1025
13.0	110	165	215	1300	1040
13.2	110	165	220	1320	1055
13.4	110	170	225	1340	1070
13.6	115	170	225	1360	1090
13.8	115	175	230	1380	1105
14.0	115	175	235	1400	1120
14.2	120	180	235	1420	1135
14.4	120	180	240	1440	1150
14.6	120	185	245	1460	1170
14.8	125	185	245	1480	1185
15.0	125	190	250	1500	1200
15.2	125	190	255	1520	1215
15.4	130	195	255	1540	1230
15.6	130	195	260	1560	1250
15.8	130	200	265	1580	1265
16.0	135	200	265	1600	1280
16.2	135	205	270	1620	1295
16.4	135	205	275	1640	1310
16.6	140	210	275	1660	1330
16.8	140	210	280	1680	1345
17.0	140	215	285	1700	1360
17.2	145	215	285	1720	1375
17.4	145	215	290	1740	1390

17.6	145	220	295	1760	1410
17.8	150	225	295	1780	1425
18.0	150	225	300	1800	1440
18.2	150	230	305	1820	1455
18.4	155	230	305	1840	1470
18.6	155	235	310	1860	1490
18.8	155	235	315	1880	1505
19.0	160	240	315	1900	1520
19.2	160	240	320	1920	1535
19.4	160	240	325	1940	1550
19.6	165	245	325	1960	1570
19.8	165	250	330	1980	1585
20.0	165	250	335	2000	1600

^a Volumes in these columns are rounded to the nearest 5 ml.

^b Give 2-hourly feeds for at least the first day. When there is little or no vomiting, moderate diarrhoea (< 5 watery stools per day), and the child finishes most feeds, change to 3-hourly feeds.

^c After a day on 3-hourly feeds, if there is no vomiting, less diarrhoea, and the child finishes most feeds, change to 4-hourly feeds.



Annex 6. How to Insert a Nasogastric Tube (NGT)

Before Insertion

- ◆ NGT feeding is usually received poorly by mothers, since it is considered invasive. Mothers should be counselled on:
 - ◆ How the tube will assist the child
 - ◆ Anticipated discomfort during insertion
 - ◆ Improved comfort after insertion
- ◆ Educate the mother on the path the NGT will take (i.e., an already existing connection from the nose, through the throat, and into the stomach). Allow the mother to express her fears, concerns, and questions to facilitate acceptance and adherence.
- ◆ This is a clean procedure; health care workers should, therefore, wash their hands thoroughly with soap prior to putting on gloves in readiness for the insertion. The child's face and torso should also be washed with soap and rinsed.
- ◆ Once correctly inserted, conduct a confirmatory test to ensure the NGT is in the stomach by:
 - ◆ Aspirating abdominal contents and testing on a litmus paper. A pH of less than 7 confirms stomach placement.
 - ◆ Pushing in air using the feeding syringe and listening to the abdomen for air sounds as you push in the air.



Feeding Using an NGT

- ◆ Emphasise the following messages with the caregiver: good hygiene practices, washing hands before feeds, and keeping utensils clean and dry.
- ◆ Assist the primary caregiver with administering the first feed; demonstrate the correct feeding position, which is the upright position.
- ◆ After attaching the feeding syringe to the NGT, milk should be poured in and allowed to flow downward freely using gravity. In cases where the free flow of milk is not achieved, raise up the feeding syringe and NGT apparatus to a higher position.
- ◆ The NGT can also be squeezed for several seconds and then released to facilitate the flow.
- ◆ ONLY when these attempts fail should a plunger be used in a slow, twisting motion. This will aid in pushing the milk downward with minimal air entry.

-
- ◆ Caregivers should be allowed to feed the child using the NGT once given a demonstration on how to feed correctly using an NGT.
 - ◆ Feeding should be done when the child is calm to avoid backflow of gastric contents; if the child is still breastfeeding, allow the child to be breastfed.
 - ◆ Change the tube if blocked. Do not plunge F-75 through the NGT; let it drip in or use gentle pressure.
 - ◆ Abdominal distension can occur with oral or NGT feeding, but it is more likely with NGT feeding. If the child develops a hard, distended abdomen with very little bowel sound, give 2 ml of a 50% solution of magnesium sulphate IM.
 - ◆ Remove the NGT when the child either takes:
 - ◆ 80% of the day's amount orally; or
 - ◆ Two consecutive feeds fully by mouth.

Exception: If a child takes two consecutive feeds fully by mouth during the night, wait until morning to remove the NGT, just in case it is needed again in the night.



Annex 7. F-100 Reference Tables

Weight of Child (kg)	Range of Volumes per 3-Hourly Feed of F-100 (8 Feeds Daily) *		Range of Volumes per 4-Hourly Feed of F-100 (6 Feeds Daily) *		Range of Daily Volumes of F-100	
	Minimum ml	Maximum ml	Minimum ml	Maximum ml	Minimum (150 ml/kg/day)	Maximum (220 ml/kg/day)
2.0	40	55	50	75	300	440
2.2	40	60	55	80	330	484
2.4	45	65	60	90	360	528
2.6	50	70	65	95	390	572
2.8	55	75	70	105	420	616
3.0	55	85	75	110	450	660
3.2	60	90	80	115	480	704
3.4	65	95	85	125	510	748
3.6	70	100	90	130	540	792
3.8	70	105	95	140	570	836
4.0	75	110	100	145	600	880
4.2	80	115	105	155	630	924
4.4	85	120	110	160	660	968
4.6	85	125	115	170	690	1012
4.8	90	130	120	175	720	1056
5.0	95	140	125	185	750	1100
5.2	100	145	130	190	780	1144
5.4	100	150	135	200	810	1188
5.6	105	155	140	205	840	1232
5.8	110	160	145	215	870	1276
6.0	115	165	150	220	900	1320
6.2	115	170	155	225	930	1364
6.4	120	175	160	235	960	1408
6.6	125	180	165	240	990	1452
6.8	130	185	170	250	1020	1496
7.0	130	195	175	255	1050	1540
7.2	135	200	180	265	1080	1584
7.4	140	205	185	270	1110	1628
7.6	145	210	190	280	1140	1672
7.8	145	215	195	285	1170	1716

8.0	150	220	200	295	1200	1760
8.2	155	225	205	300	1230	1804
8.4	160	230	210	310	1260	1848
8.6	160	235	215	315	1290	1892
8.8	165	240	220	325	1320	1936
9.0	170	250	225	330	1350	1980
9.2	175	255	230	335	1380	2024
9.4	175	260	235	345	1410	2068
9.6	180	265	240	350	1440	2112
9.8	185	270	245	360	1470	2156
10.0	190	275	250	365	1500	2200
10.2	190	280	255	375	1530	2244
10.4	195	285	260	380	1560	2288
10.6	200	290	265	390	1590	2332
10.8	205	295	270	395	1620	2376
11.0	205	305	275	405	1650	2420
11.2	210	310	280	410	1680	2464
11.4	215	315	285	420	1710	2508
11.6	220	320	290	425	1740	2552
11.8	220	325	295	435	1770	2596
12.0	225	330	300	440	1800	2640
12.2	230	335	305	445	1830	2684
12.4	235	340	310	455	1860	2728
12.6	235	345	315	460	1890	2772
12.8	240	350	320	470	1920	2816
13.0	245	360	325	475	1950	2860
13.2	250	365	330	485	1980	2904
13.4	250	370	335	490	2010	2948
13.6	255	375	340	500	2040	2992
13.8	260	380	345	505	2070	3036
14.0	265	385	350	515	2100	3080
14.2	265	390	355	520	2130	3124
14.4	270	395	360	530	2160	3168
14.6	275	400	365	535	2190	3212
14.8	280	405	370	545	2220	3256
15.0	280	415	375	550	2250	3300
15.2	285	420	380	555	2280	3344
15.4	290	425	385	565	2310	3388
15.6	295	430	390	570	2340	3432
15.8	295	435	395	580	2370	3476
16.0	300	440	400	585	2400	3520


16.2	305	445	405	595	2430	3564
16.4	310	450	410	600	2460	3608
16.6	310	455	415	610	2490	3652
16.8	315	460	420	615	2520	3696
17.0	320	470	425	625	2550	3740
17.2	325	475	430	630	2580	3784
17.4	325	480	435	640	2610	3828
17.6	330	485	440	645	2640	3872
17.8	335	490	445	655	2670	3916
18.0	340	495	450	660	2700	3960
18.2	340	500	455	665	2730	4004
18.4	345	505	460	675	2760	4048
18.6	350	510	465	680	2790	4092
18.8	355	515	470	690	2820	4136
19.0	355	525	475	695	2850	4180
19.2	360	530	480	705	2880	4224
19.4	365	535	485	710	2910	4268
19.6	370	540	490	720	2940	4312
19.8	370	545	495	725	2970	4356
20.0	375	550	500	735	3000	4400



Annex 8. Toys for Children



Ring on a string (from 6 months)

Thread cotton reels and other small objects (e.g. cut from the neck of plastic bottles) on to a string. Tie the string in a ring, leaving a long piece of string hanging.



In-and-out toy (from 9 months)

Any plastic or cardboard container and small objects (not small enough to be swallowed).




Blocks (from 9 months)

Small blocks of wood. Smooth the surfaces with sandpaper and paint in bright colours, if possible.


Push-along toy (from 12 months)

Make a hole in the centre of the base and lid of a cylindrical-shaped tin. Thread a piece of wire (about 60 cm long) through each hole and tie the ends inside the tin. Put some metal bottle tops inside the tin and close the lid.



Pull-along toy (from 12 months)

As above, except that string is used instead of wire.



Doll (from 12 months)

Cut out two doll shapes from a piece of cloth and sew the edges together, leaving a small opening. Turn the doll inside-out and stuff with scraps of materials. Stitch up the opening and sew or draw a face on the doll.

Rattle (from 12 months)

Cut long strips of plastic from coloured plastic bottles. Place them in a small transparent plastic bottle and glue the top on firmly.



Drum (from 12 months)


Any tin with a tightly fitting lid.

Mirror (from 18 months)

A tin lid with no sharp edges.


Posting bottle (from 12 months)

A large transparent plastic bottle with a small neck and small long objects that fit through the neck (not small enough to be swallowed).



Stacking bottle tops (from 12 months)

Cut at least three identical round plastic bottles in half and stack them.




Nesting toys (from 9 months)

Cut off the bottom of two bottles of identical shape, but different size. The smaller bottle should be placed inside the larger bottle.



Puzzle (from 18 months)

Draw a figure (e.g. a doll) using a crayon on a square- or rectangular-shaped piece of cardboard. Cut the figure in half or quarters.



Book (from 18 months)

Cut out three rectangular-shaped pieces of the same size from a cardboard box. Glue or draw a picture on both sides of each piece. Make two holes down one side of each piece and thread string through to make a book.



WHO 97402



Annex 9. F-100-Diluted Reference Table for Infant 0-6 Months Old

Reference Table for Maintenance Amounts of F-100-Diluted to Give to an Individual Breastfed Infant per Feed	
Bodyweight (kg)	F-100-Diluted (ml per Feed) (Assumes 8 Feeds per Day, Given 3-Hourly)
1.3	25
1.3–1.5	30
1.6–1.7	35
1.8–2.1	40
2.2–2.4	45
2.5–2.7	50
2.8–2.9	55
3.0–3.4	60
3.5–3.9	65
4.0–4.4	70

Reference Table for Amounts of F-100-Diluted or F-75 (with Bilateral Pitting Oedema) to Give to Non-Breastfed Infants in the Stabilisation Phase	
Bodyweight (kg)	F-100-Diluted or F-75 (ml per Feed), 8 Feeds per Day, No Breastfeeding (3-Hourly Feeds)
≤ 1.6	30
1.6–1.8	35
1.9–2.1	40
2.2–2.4	45
2.5–2.7	50
2.8–2.9	55
3.0–3.4	60
3.5–3.9	65
4.0–4.4	70


Reference Table for Amounts of F-100-Diluted to Give to Non-Breastfed Infants 0–6 Months in the Transition Phase

Bodyweight (kg)	F-100-Diluted (ml per Feed), 8 Feeds per Day, No Breastfeeding (3-Hourly Feeds)
≤ 1.6	45
1.6–1.8	53
1.9–2.1	60
2.2–2.4	68
2.5–2.7	75
2.8–2.9	83
3.0–3.4	90
3.5–3.9	96
4.0–4.4	105

Reference Table for Amounts of F-100-Diluted to Give to Non-Breastfed Infants 0–6 Months in the Rehabilitation Phase

Bodyweight (kg)	F-100-Diluted (ml per Feed), 6 to 8 Feeds per Day, No Breastfeeding
≤ 1.6	60
1.6–1.8	70
1.9–2.1	80
2.2–2.4	90
2.5–2.7	100
2.8–2.9	110
3.0–3.4	120
3.5–3.9	130
4.0–4.4	140

Annex 10. Decision Tree on the Management of MAM



MAM prevention and treatment interventions should be linked with SAM treatment, IYCF and SBCC, WASH and other health and nutrition interventions. Households of MAM patients should also be referred for PSNP throughout the year.

MAM treatment with TSFP should be integrated with routine health services and OTP where possible.



Annex 11. Specialised Nutritious Foods

Objective	Treatment of Severe Acute Malnutrition (SAM)		Treatment of Moderate Acute Malnutrition (MAM)	
Generic Term	Ready-to-Use Therapeutic Foods (RUTF)		Ready-to-Use Supplementary Foods (RUSF)	
Purpose	RUTF given during treatment of uncomplicated SAM with continued breastfeeding		RUSF given to treat moderate acute malnutrition with continued breastfeeding	
Target Group	6–59 months		6–59 months, others including pregnant and lactating women and older people (≥ 60 years)	
Energy and Nutrient/ Ration or Dose	500 kcal 12.5g protein 31.9g fat		510 kcal 13g protein (10% energy) 31g fat (55% energy) Meets WHO nutrient content guidance for MAM foods.	
Packaging	Packet = 92g		Packet = 100g	
Shelf Life	24 months		24 months	
Ration or Dose	According to weight: 200 kcal/kg/day		1 packet/day 100g/day	
			200g/day	

Adapted from: Global Nutrition Cluster. 2017. Moderate Acute Malnutrition: A Decision Tool for Emergencies. Available at: <http://nutritioncluster.net/?get=002086%7C2014/07/MAM-Decision-Tool-final-June-2014-corrected.pdf>.

Annex 12. Nutrient Requirements for PLHIV

Nutrient needs depend on age; physical changes, such as pregnancy and breastfeeding; and level of activity. For PLHIV, energy requirements are influenced by severity of disease state. The nutrient requirements of various groups of people are shown below.

Age group	Healthy	HIV-infected		
		Asymptomatic	Symptomatic	Severely undernourished
Children		10% more energy	20% more energy	50%–100% more energy
6–11 months	690	760	830	150–220 kcal/kg of body weight/day
12–23 months	900	990	1080	150–220 kcal/kg of body weight/day
2–5 years old	1260	1390	1510	150–220 kcal/kg of body weight/day
6–9 years old	1650	1815	1980	75–100 kcal/kg of body weight/day
10–14 years old	2,020	2,220	2,420	60–90 kcal/kg of body weight/day
Adults ≥ 18 years				
Non-pregnant and lactating	2,000–2,580	10% more energy	20% more energy	
Pregnant and lactating women	2,460–2,570*	(210–258 more kcal)	(420 more kcal)	

* The requirements for adults also apply to pregnant and lactating women, in addition to the usual extra requirements for pregnancy and lactation.

Source: Adapted from: WHO. 2009.

Protein

Protein should constitute 12%–15% of dietary intake (50–80 g/day or 1 g/kg of ideal body weight).

According to WHO, there is no evidence that PLHIV have different protein requirements than healthy HIV-negative people.

Fat

Fat/oil intake should not be more than 35% of total energy needs. People with HIV and/or TB should consume the same percentage of energy from fat as healthy people.

Vitamins and Minerals

Eating a varied diet is the best way to ensure adequate intake of vitamins and minerals. Where dietary intake of vitamins and minerals may not be enough to correct nutritional deficiencies, or the recommended intakes cannot be achieved, high-risk groups, such as pregnant and lactating women, may need multiple micronutrient supplements.

Annex 13. Essential Commodities for the Management of SAM

Stabilisation Centre Essential Commodities for the Management of SAM with Medical Complications

Item or Product	Unit	Minimum Quantity Per Facility
<i>Medical Supplies and Equipment</i>		
Adhesive plaster	Packet	1
Pediatric cannula	Packet	3
Feeding syringe (60ml)	Packet	1
Syringe with needle (5ml)	Packet	8
Nasogastric tube (size 6)	Packet	2
Nasogastric tube (size 8)	Packet	2
Nasogastric tube (size 12)	Packet	2
Thermometer	Piece	1
<i>Medicine</i>		
Mebendazole or Albendazole	Tablet	Based on expected caseload.
Amoxicillin	Tablet	Based on expected caseload.
Ampicillin IV	Vial	Based on expected caseload.
Water for injection (5ml)	Vial	Based on expected caseload.
Gentamicin IV	Vial	Based on expected caseload.
Paracetamol	Bottle	Based on expected caseload.
Ceftriaxone IV	Vial	Based on expected caseload.
Ciprofloxacin	Tablet	Based on expected caseload.
Cloxacillin	Vial	Based on expected caseload.
Glucose solution IV	Packet	Based on expected caseload.
Vitamin A	Tablet	Based on expected caseload.
ReSoMal	Packet	Based on expected caseload.
<i>Anthropometric Equipment</i>		
Weighing scale	Piece	1
MUAC tapes (child)	Piece	4
Length/height board	Piece	1
<i>Specialised Nutritious Foods</i>		
F-75	Tin	Based on expected caseload.
F-100	Tin	Based on expected caseload.
RUTF	Packet	Based on expected caseload.
<i>Monitoring, Reporting and Reference Materials</i>		
Stabilisation Centre Multi-chart	Piece	Based on expected caseload.

Referral Slip	Piece	20
Monthly Statistics Report for SAM and MAM	Piece	1
Monthly Supplies Report for SAM and MAM	Piece	1
Registration Book for SAM	Piece	1
WFH/WFL Reference Tables	Piece	1
Quick Reference Guide	Piece	1
Guidelines for the Management of Acute Malnutrition	Piece	1
IEC Materials	Set	1
Food Preparation and Other Materials		
Cups/tumblers	Piece	10
Spoons	Piece	10
Water jug with lid	Piece	2
Wooden pallets	Piece	2

OTP for the Management of SAM without Medical Complications

Item or Product	Unit	Minimum Quantity Per Facility
<i>Medical Supplies and Equipment</i>		
Thermometer	Piece	1
<i>Medicine</i>		
Mebendazole or Albendazole	Tablet	Based on expected caseload.
Amoxicillin	Tablet	Based on expected caseload.
<i>Anthropometric Equipment</i>		
Weighing scale	Piece	1
MUAC tapes (child)	Piece	4
<i>Specialized Nutritious Foods</i>		
RUTF	Packet	Based on expected caseload.
Monitoring, Reporting and Reference Materials		
OTP Treatment and Follow-up Cards	Piece	Based on expected caseload.
Referral Slip	Piece	20
Registration Book for SAM	Piece	1
Monthly Statistics Report for SAM and MAM	Piece	1
Monthly Supplies Report for SAM and MAM	Piece	1
Quick Reference Guide	Piece	1
Guidelines for the Management of Acute Malnutrition	Piece	1
IEC Materials	Set	1
Food Preparation and Other Materials		
Cups/tumblers	Piece	10
Spoons	Piece	10
Water jug with lid	Piece	2
Wooden pallets	Piece	2

Annex 14. Essential Commodities for the Management of MAM

Item or Product	Unit	Minimum Quantity Per Facility
<i>Medicine</i>		
Mebendazole or Albendazole	Tablet	Based on expected caseload
Vitamin A	Tablet	Based on expected caseload
<i>Anthropometric Equipment</i>		
Weighing scale	Piece	1 to 2 depending on facility caseload
MUAC tapes (child)	Piece	4
MUAC tapes (adult)	Piece	4
<i>Specialised Nutritious Foods</i>		
RUSF or FBF	Packet	Based on expected caseload
<i>Monitoring, Reporting and Reference Materials</i>		
TSFP Treatment and Follow-up Card	Piece	Based on expected caseload
TSFP Identification Card	Piece	Based on expected caseload
Referral Slip	Piece	20
Registration Book for MAM—Children 6–59 months	Piece	1
Registration Book for MAM—PLW	Piece	1
Monthly Statistics Report for SAM and MAM	Piece	1
Monthly Supplies Report for SAM and MAM	Piece	1
Quick Reference Guide	Piece	1
Guidelines for the Management of Acute Malnutrition	Piece	1
IEC Materials	Set	1
<i>Food Preparation and Other Materials</i>		
Cooking Demonstration Equipment	Piece	1
Cups/tumblers	Piece	20
Spoons	Piece	20
Water jug with lid	Piece	2
Wooden pallets	Piece	5



Annex 17. OTP Treatment and Follow-up Card

ADMISSION DETAILS: OTP Treatment Card

Full Name					Registration N°	/ /			
Age (months)	Sex		M	F	Date of admission				
Mother's Name					Reg. N°				
Region	Woreda			Kebele					
OTP Site					Distance to the house (hours)				
Referred by	Community Volunteer			Other (neighbours etc.)			Self-referred		
Admission (circle)	New Case	Relapse	Returned Defaulter	From Routine or Community Health Days	From Stabilisation Centre	Stabilisation Centre Refusal			
2nd Admission (relapse)	No	Yes	Additional information						
Admission Anthropometry									
lateral pitting oedema	+ ++ +++								
MUAC (cm)	Weight (kg)		Height/length (cm)		Weight for height/length (WFL)				
Admission criteria	lateral pitting	MUAC	Weight for height	Other:					
History									
Diarrhoea	yes	no	# Stools/day		1-3	4-5	>5		
Vomiting	yes	no	Passing urine		yes	no			
Cough	yes	no	If bilateral pitting oedema, how long swollen?						
Appetite	good	poor	none	Breastfeeding status (children 6 to 23 months)		yes	no		
Additional information									
Physical Examination									
Respir. rate (#/min)	<30	30 – 39	40 - 49	50+	Chest Indrawing		yes	no	
Temperature °C					Conjunctiva		normal	pale	
Eyes	normal	sunken	discharge		Dehydration	None	moderate	severe	
Ears	normal	discharge		Mouth		Normal	sores	candida	
Enlarged lymph nodes	none	neck	axilla	groin	Hands & feet		normal	cold	
Skin changes	none	scabies	peeling	ulcers / abscesses		Disability		yes	no
Additional information									
Routine Medications									
1st VISIT:	Drug	date	Dosage		Drug	Date	Dosage		
	Amoxicillin								
	Malaria treatment	yes	no	date	Measles Immunisation	No	Yes	date:	
	2nd VISIT:	drug	date	Dosage	Fully immunised	No	Yes		
	Mebendazole/Albendazole								
Other Medication									
	Drug	Date	Dosage		Drug	Date	Dosage		
Transfer-in and Transfer-out during treatment for SAM									
Transfer In				Transfer Out					
Location	Date	Reg No of other Facility		Reason	Location	Date	Reg No.		
Home Visit									
Date	Reason for Home Visit			Date of Home Visit	Findings				

FOLLOW UP: OTP

NAME	Registration N°																
Week	ADM (=0)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Date																	
Anthropometry																	
Bilateral Pitting Oedema (+ ++ +++)																	
MUAC (cm)																	
Weight (kg)				*		*											
Weight loss * (Y/N)																	
Height/length (cm)																	
Weight for Height/length																	
* If below admission weight on week 3 refer for home visit; If no weight gain by week 5 refer to SC																	
History																	
Diarrhoea (# days)																	
Vomiting (# days)																	
Fever (# days)																	
Cough (# days)																	
Physical examination																	
Temperature (°C)																	
Respiratory rate (# /min)																	
Dehydrated (Y/N)																	
Anaemia / palmar pallor																	
Skin infection (Y/N)																	
Appetite test / feeding																	
RUTF test (Passed/Failed)																	
RUTF (# units given)																	
Breastfeeding																	
Action / follow up																	
ACTION NEEDED (Y/N)																	
Other medication (see Name of examiner)																	
VISIT OUTCOME																	
OK=Continue A=Absent DF= Defaulter (Absent for 2 consecutive visits and confirmed by a home visit) T=Transfer RR=Refused referral C=Cured NR=Non-responder HV= Home visit D=Died																	
Action taken during follow-up (include date)																	
Name of outreach worker:																	



Annex 19. TSFP Treatment and Follow-up Card for Children 6-59 Months

TSFP Treatment and Follow-up Card for Children 6-59 Months of Age

Mother's/Guardian's Name:		Registration No:	
Child's Name:		Sex (M/F):	
Date of Admission:		Age (months):	
Health Post:		Village/Gott:	

Follow-up Visit	Admission	1	2	3	4	5	6	7	8
Date									
Anthropometry									
Weight (kg)			*		*				
MUAC (cm)									
**Bilateral Pitting Odema (0, +, ++, +++)									
* If below admission weight on 4th week in TSFP conduct home visit, if no weight gain on 8th week in TSFP refer for further medical assessment. **If a child has bilateral pitting oedema, they should be referred to the OTP and treated according to the Action Protocol.									
Medical History									
Diarrhoea (# days)									
Vomiting (# days)									
Fever (# days)									
Cough (# days)									
Routine Medication									
Vitamin A									
Mebendazole/ Albendazole									
Type of Specialised Nutritious Food									
Quantity/Ration size (No. of sachets/ packets)									
Signature/ Thumb print									
Action & Visit Outcome									
Action Needed (Y/N)									
Visit Outcome									
CODES for Visit Outcome: OK=Continue A=Absent DF=Absent for 2 consecutive visits and confirmed by a home visit T=Transfer RR=Refused referral C=Cured NR=Non-responder HV= Home visit D=Died									

Annex 20. TSFP Treatment and Follow-up Card for Pregnant and Lactating Women

TSFP Treatment and Follow-up Card for Pregnant and Lactating Women

Full Name:								
Date of Admission:								
Pregnant Woman: Expected Delivery Date:								
Lactating Woman: Date the baby reaches 6 months:								
	Registration No:							
	Kebele:							
	Village/Gott:							
	Health Post:							

	1	2	3	4	5	6	7	8
Follow-up Visit	Admission							
Date								
Weight (Kg)								
MUAC (cm)								
Mebendazole/ Albendazole								
Type of Specialised Nutritious Food								
Quantity/Ration size (No. of sachets/ packets)								
Signature/ Thumb print								
Visit Outcome								

CODES for Visit Outcome: OK=Continue A=Absent DF=Absent for 2 consecutive visits and confirmed by a home visit T=Transfer RR=refused referral C=Cured NR=Non-responder HV=Home visit D=Died Other: pregnant woman who has delivered or lactating woman whose infant is ≥ 6 months



Annex 21. TSFP Identification Card for Admitted Patients

TSFP Identification Card for Admitted Patients.

Name:		Registration No:	
Mother's/Guardian's Name:		Date of Admission:	
Sex (M/F):		Age(in Months):	
Pregnant Woman: Expected Delivery Date:		Village/Gott:	
Lactating Woman: Date the infant reaches 6 months:		Health Post:	

TSFP Identification Card for Admitted Patients.

Name:		Registration No:	
Mother's/Guardian's Name:		Date of Admission:	
Sex (M/F):		Age(in Months):	
Pregnant Woman: Expected Delivery Date:		Village/Gott:	
Lactating Woman: Date the infant reaches 6 months:		Health Post:	

TSFP Identification Card for Admitted Patients.

Name:		Registration No:	
Mother's/Guardian's Name:		Date of Admission:	
Sex (M/F):		Age(in Months):	
Pregnant Woman: Expected Delivery Date:		Village/Gott:	
Lactating Woman: Date the infant reaches 6 months:		Health Post:	

TSFP Identification Card for Admitted Patients.

Name:		Registration No:	
Mother's/Guardian's Name:		Date of Admission:	
Sex (M/F):		Age(in Months):	
Pregnant Woman: Expected Delivery Date:		Village/Gott:	
Lactating Woman: Date the infant reaches 6 months:		Health Post:	



Annex 22. Referral Slip for the Management of Acute Malnutrition

Full Name:		Registration No:	
Caregiver/Mother's Name:		Sex (M/F):	
Woreda:		Age:	
Kebele:		Referred From:	
Gott:		Date of Referral:	
REFERRAL DATA:			
	On Admission	During Referral	
MUAC:			
Weight:			
Height:			
WFL/WFH:			
INFANTS 0–6 MONTHS OF AGE:			
Is infant exclusively breastfeeding? (circle) Y N			
REFERRAL NOTES:			
Referral from OTP to SC or Further Medical Investigations:			
Reason for Referral	Circle as Appropriate		
Bilateral pitting oedema (+++)	Y	N	
Severe wasting with any bilateral pitting oedema	Y	N	
Severe wasting or bilateral pitting oedema (or ++) with medical complications	Y	N	
Poor appetite and deteriorating nutrition and/or medical condition based on the action protocol.	Y	N	
Referred for further medical investigations.	Y	N	
Referral from SC to OTP:			
Reason for Referral	Circle as Appropriate		
Stabilised medical complications	Y	N	
Treatment or Medications Given:			
Treatment/medications	Specify		
Antibiotic			
Malaria Treatment			
Mebendazole/Albendazole			
Measles Immunisation			
Other medications.....			
Comments:			

Annex 26. Monthly Statistics Report for SAM and MAM

Region: _____
 Zone: _____
 Woreda: _____
 Type of Facility: _____
 Name of Facility: _____
 Report Prepared by: _____
 Month/Year of Reporting: _____
 Opening Date: _____

SEVERE ACUTE MALNUTRITION (SAM)

Service	Age	Total at the beginning of the month (A)	New Admissions		Re-admissions		Transfers-in from OTP, SC or other facility (E)	Total Admissions [(F)=B+C+D+E]	Cured (G)	Died (H)	Defaulter (I)	Non-responder (J)	Stabilised to OTP (K)	Transfers-out to SC or other facility (L)	Total Discharged [(M)=G+H+I+L]	Total at the end of the month [(N)=A+M]
			MUAC or WFL/WFL (B)	Bilateral pitting oedema (C)	Relapse (D)	Returned Defaulters (E)										
	6-59 months															
	Other, e.g. Older people															
	Total															
	< 6 months															
	6-59 months															
	Other, e.g. Older people															
	Total															
Stabilisation Centre (SC)																
TARGET (Sphere Standards) % >75% % <10% % <15% % <15% %																

MODERATE ACUTE MALNUTRITION (MAM)

Service	Age	Total at the beginning of the month (A)	New Admissions		Re-admissions		Transfers-in from other facility (E)	Total Admissions [(F)=B+C+D+E]	Cured (G)	Died (H)	Defaulter (I)	Non-responder (J)	Transfers-out to other facility (K)	Others, pregnant and lactating or infant > 6 months (L)	Total Discharged [(M)=G+H+I+L]	Total at the end of the month [(N)=A+M]
			MUAC or WFL/WFL (B)	Bilateral pitting oedema (C)	Relapse (D)	Returned Defaulters (E)										
Targeted Supplementary Feeding Programme (TSFP) for Children 6-59 months and other	6-59 months															
	Other, e.g. Older people															
	Total															
Pregnant																
Lactating																
TSFP for Pregnant and Lactating Women	Total															
TARGET (Sphere Standards) % >75% % <3% % <15% %																

Report Completed By: _____ Date: _____
 Report Approved By: _____ Date: _____



Annex 27. Monthly Supplies Report for SAM and MAM

Region: _____
 Zone: _____
 Woreda: _____
 Type of Facility: _____
 Name of Facility: _____
 Report Prepared by: _____
 Month/Year of Reporting: _____
 Opening Date: _____

Supply Management for SAM

Item or Product	Unit	Stock balance at the beginning of the month (A)	Received in the month (B)	Issued in the month (C)	Losses or Adjustments (D)	Stock balance at the end of the month [(E) = A+B-C-D]	Request for Supplies			Remarks
							Quantity needed	Unit	Period of request in months	
RUTF	Sachets									
F-75	Tins									
F-100	Tins									
ReSoMal	Sachets									
SC Kits	Kits									
Other, specify: _____										

Supply Management for MAM

Item or Product	Unit	Stock balance at the beginning of the month (A)	Received in the month (B)	Issued in the month (C)	Losses or Adjustments (D)	Stock balance at the end of the month [(E) = A+B-C-D]	Request for Supplies			Remarks
							Quantity needed	Unit	Period of request in months	
RUSF	Sachets									
Fortified Blended Food (FBF), e.g. Super Cereal Plus	Packets									
Other, specify: _____										

Report Completed By: _____ Date: _____
 Report Approved By: _____ Date: _____



Ministry of Health

National Guideline for the Management of Acute Malnutrition in Ethiopia
May 2019